

Existence and Uniqueness of the Measure of Maximal Entropy for the Teichmüller Flow on the Moduli Space of Abelian Differentials.

Alexander I. Bufetov* and Boris M. Gurevich†

1 Introduction

The Teichmüller geodesic flow $\{g_t\}$, first studied by H. Masur [15] and W. Veech [21], acts on the moduli space of Riemann surfaces endowed with a holomorphic differential. More precisely, let S be a closed surface of genus $g \geq 2$. One introduces on S a complex structure σ and a holomorphic differential ω . The pair (σ, ω) is considered to be equivalent to another pair of the same nature (σ_1, ω_1) if there is a diffeomorphism of S sending (σ, ω) to (σ_1, ω_1) . The moduli space $\mathcal{M}(g)$ consists of the equivalence classes, and the flow $\{g_t\}$ on $\mathcal{M}(g)$ is induced by the action on the pairs (σ, ω) defined by the formula $g_t(\sigma, \omega) = (\sigma', \omega')$, where $\omega' = e^t \Re(\omega) + ie^{-t} \Im(\omega)$, while the complex structure σ' is determined by the requirement that ω' be holomorphic. If (σ, ω) and (σ', ω') are equivalent, then the differentials ω and ω' have the same orders of zeros and the same area. Therefore, these orders and area are well-defined on $\mathcal{M}(g)$. Moreover, they are preserved by the Teichmüller flow $\{g_t\}$. Take an arbitrary non-ordered collection $\kappa = (k_1, \dots, k_r)$ with $k_i \in \mathbb{N}$, $k_1 + \dots + k_r = 2g - 2$, and denote by $\mathcal{M}_\kappa$ the subspace of $\mathcal{M}(g)$ corresponding to the differentials of area 1 (i.e., $(i/2) \int \omega \wedge \bar{\omega} = 1$) with orders of zeros k_i , $i = 1, \dots, r$; $\mathcal{M}_\kappa$ is said to be a *stratum* in $\mathcal{M}(g)$. Each stratum

*Department of Mathematics, Rice University, and the Steklov Institute of Mathematics, Russian Academy of Sciences.

†Department of Mechanics and Mathematics, Moscow State University, and the Institute for Information Transmission Problems, Russian Academy of Sciences.

is a $\{g_t\}$ -invariant set, and there is a natural $\{g_t\}$ -invariant measure on $\mathcal{M}_\kappa$; this measure is finite [15], [21].

The space $\mathcal{M}_\kappa$ also admits a natural topological structure, in which it is in general non-connected. The number of connected components is no more than 3 and depends on κ (see [13] for details), each of them is $\{g_t\}$ -invariant.

We fix an arbitrary closed component $\mathcal{H}$ and denote by μ_κ the normalized restriction to $\mathcal{H}$ of the above-mentioned $\{g_t\}$ -invariant measure.

Veech [23] showed that $\{g_t\}$ with respect to the measure μ_κ is a Kolmogorov flow with entropy given by the formula

$$h_{\mu_\kappa}(\{g_t\}) = 2g - 1 + r. \quad (1.1)$$

Our aim is to establish the following

Theorem 1.1. *The measure μ_κ is the unique measure of maximal entropy for the flow $\{g_t\}$ on $\mathcal{H}$.*

The proof of this theorem is based on the representation of the flow $\{g_t\}$ as a suspension flow over a countable alphabet topological Markov shift. The reasoning proceeds in two steps. We begin with sufficient conditions for an invariant measure of the above suspension flow to be a measure with maximal entropy. These conditions are contained in Theorem 2.2 stated in Subsection 2.1. After stating the theorem we outline its proof. This proof is close in spirit to thermodynamic formalism for countable alphabet topological Markov shifts [12], [20], [19]. In particular, we use a uniqueness theorem by Buzzi and Sarig [6] for an equilibrium measure. An application of thermodynamic formalism to another smooth dynamical system with non-compact phase space, the geodesic flow on the modular surface, can be found in [11].

Subsections 2.2–2.5 are devoted to the proof of Theorem 2.2 in detail. In the rest of the paper (Sections 3–5) we deduce Theorem 1.1 from Theorem 2.2, and there (at the beginning of Section 3), as before, we start from a sketch of the subsequent reasoning.

The following observation lies at the centre of our argument in this part of the proof. The Teichmüller flow admits infinitely smooth stable and unstable foliations, with respect to which it is “measurably Anosov” in the sense of Veech [23] and Forni [8]. The Masur-Veech measure μ_κ induces globally defined sigma-finite measures on unstable leaves and these measures are uniformly expanded by the flow. In other words, the Masur-Veech measure has

the Margulis [14] uniform expansion property on unstable leaves. Informally, Proposition 4.8 expresses the Margulis property in terms of the symbolic representation of the Teichmüller flow.

Let us remark that to establish Theorem 1.1 we need only the special case of Theorem 2.2 dealing with the countable alphabet topological Bernoulli shift. But the proof for this case would be only a little easier than in the general one. The main results of this paper are stated without proof in [5].

2 Suspension flows

Let G be an Abelian group (in what follows only $G = \mathbb{Z}$ or $G = \mathbb{R}$ will appear), and let $\{T_g, g \in G\}$ be an action of G by measurable transformations of a metrizable topological space X endowed with its Borel σ -algebra $\mathcal{B}$. Two actions, $\{X, T_g\} = \{T_g, g \in G\}$ on $(X, \mathcal{B})$ and $\{X', T'_g\} = \{T'_g, g \in G\}$ on $(X', \mathcal{B}')$, are called *isomorphic* if there is a one-to-one epimorphic bimeasurable map $\Phi : X \rightarrow X'$ such that $T'_g \circ \Phi = \Phi \circ T_g$ for all $g \in G$. If Φ is not necessarily epimorphic, we say that $\{X, T_g\}$ is *embedded* into $\{X', T'_g\}$.

Consider also an action $\{X, T_g\}$ together with a $\{T_g\}$ -invariant Borel probability measure μ on X . Denote such an object by $\{X, T_g; \mu\}$. We say that $\{X, T_g; \mu\}$ and $\{X', T'_g; \mu'\}$ are isomorphic if there are sets $X_1 \in \mathcal{B}$, $X'_1 \in \mathcal{B}'$ invariant with respect to all T_g and all T'_g respectively such that $\mu(X_1) = \mu'(X'_1) = 1$ and the restrictions $\{X_1, T_g|_{X_1}\}$ and $\{X'_1, T'_g|_{X'_1}\}$ are isomorphic in the above sense.

If $G = \mathbb{Z}$, the corresponding action will be denoted by $\{X, T_n\}$. If $G = \mathbb{R}$, we write $\{X, T_t\}$ or $\{T_t\}$ (or just T_t , S_t , etc. when it cannot cause confusion). In the former case $T_n = T^n$ where T is a bimeasurable one-to-one transformation of $(X, \mathcal{B})$ called an *automorphism*. In the latter case the action is called a *flow*. We keep the same terminology for actions considered together with their invariant measures. In this paper we mostly deal with flows that can be defined as follows. Let T be an automorphism of $(X, \mathcal{B})$ and $f : X \rightarrow [c, \infty)$, $c > 0$, be a measurable function. Consider the direct product $X \times \mathbb{R}_+$ and its subspace $X_f = \{(x, u) : x \in X, 0 \leq u < f(x)\}$. For $t \geq 0$ and every point $\tilde{x} = (x, u) \in X_f$, we set $S_t \tilde{x} = (x, u + t)$ if $u + t < f(x)$, and $S_t \tilde{x} = (T^n x, u + t - \sum_{i=0}^{n-1} f(T^i x))$, where n is such that $\sum_{i=0}^{n-1} f(T^i x) \leq u + t < \sum_{i=0}^n f(T^i x)$. For $t < 0$ we set $S_t = (S_{-t})^{-1}$ and thus obtain a flow $\{X_f, S_t\}$. For this flow, we shall also use the notation (T, f) and call it the *suspension flow* constructed by T and the *roof function* f .

Denote by $\mathcal{M}_{T,f}$ the set of all T -invariant Borel probability measures μ on X with $\mu(f) < \infty$ (here and in the sequel, $\mu(f) := \int f d\mu$). Every (T, f) -invariant Borel probability measure μ_f on X_f is induced by a measure $\mu \in \mathcal{M}_{T,f}$. Namely,

$$\mu_f = (\mu(f))^{-1}(\mu \times \lambda)|_{X_f},$$

where λ is the Lebesgue measure on $\mathbb{R}_+$. We will refer to μ_f as the *f-lifting* of μ and for brevity write $(T, f; \mu_f)$ instead of $((T, f); \mu_f)$. The entropy $h(T, f; \mu_f)$ of the flow (T, f) with respect of the measure μ_f is given by Abramov's formula

$$h(T, f; \mu_f) = h(T; \mu)/\mu(f), \quad (2.1)$$

where $h(T; \mu)$ is the entropy of the automorphism T with respect to the measure μ . We define the *topological entropy* of (T, f) by

$$h_{\text{top}}(T, f) = \sup_{\mu \in \mathcal{M}_{T,f}} h(T, f; \mu_f). \quad (2.2)$$

This terminology is justified by the following well-known fact: if X is a compact space, T is a homeomorphism of X , and f is continuous, then the right-hand side of (2.2) is indeed the topological entropy of the suspension flow (T, f) .

We refer to every $\mu \in \mathcal{M}_{T,f}$ at which the supremum in (2.2) is achieved as to a *measure of maximal entropy* for (T, f) .

2.1 Suspension flows over Markov shifts

In the specific case we will deal with, (X, T) is a countable alphabet topological Markov shift, i.e., X is the set of infinite two-sided paths of a directed graph $\Gamma = (V, E)$ with vertex set V and edge set $E \subseteq V \times V$, and T is the shift transformation: $(Tx)_i = x_{i+1}$ for each $x = (x_i, i \in \mathbb{Z}) \in X$. In other words, X consists of all sequences $x \in V^{\mathbb{Z}}$ such that $B_{x_i, x_{i+1}} = 1$, where $B = B(\Gamma)$ be the incidence matrix of the graph Γ . The vertices $v \in V$ will also be called letters.

In the sequel we assume that Γ is connected. If Γ is the complete graph, i.e., $E = V \times V$, we have the *topological Bernoulli shift with alphabet V*.

We introduce the discrete topology on V , the product topology on $V^{\mathbb{Z}}$, and the induced topology on $X \subset V^{\mathbb{Z}}$. The map T is clearly a homeomorphism of X . We shall refer to every finite path of Γ , i.e., a sequence $w = (v_1, \dots, v_k) \in V^k$ such that (v_i, v_{i+1}) , $i = 1, \dots, k-1$, as a *word* and sometimes say that

this word is from v_1 to v_k . Denote the set of all words (including the empty word) by $W(\Gamma)$.

Let $w = (v_1, \dots, v_k)$, $w' = (v'_1, \dots, v'_l)$ be two words. The concatenation $ww' := (v_1, \dots, v_k, v'_1, \dots, v'_l)$ is also a word if $(v_k, v'_1) \in E$. We say that w contains w' (or w' is a subword of w) if $v'_1 = v_i, \dots, v'_l = v_{i+l-1}$ for some i , $1 \leq i \leq k - l + 1$. In the special case that $i = 1$, we call w' a *prefix* of w . Let $w = (v_1, \dots, v_n)$, $n \geq 2$, be a word and $w' = (v_1, \dots, v_l)$, $l \leq n$, be a prefix of w . We call w' a *simple prefix* of w if there is no k , $2 \leq k \leq l$, such that $(v_1, \dots, v_{n-k+1}) = (v_k, \dots, v_n)$. If w is a simple prefix of itself, then w is called a *simple word*. If a simple word is a prefix of another word, it is clearly a simple prefix.

Remark 2.1. Every word $w = (v_1, \dots, v_n)$ is certainly the concatenation of the single-letter words v_i , so we will also write $w = v_1 \dots v_n$.

To every word w we assign the cylinder $C_w = \{x \in X : (x_0, \dots, x_{|w|-1}) = w\}$, where $|w|$ is the length of w , i.e., the number of symbols in w .

For a function $f : X \rightarrow \mathbb{R}$, we set

$$\text{var}_n(f) = \sup\{|f(x) - f(y)| : x_i = y_i \text{ when } |i| \leq n\}, \quad n \in \mathbb{N}.$$

We say that f has *summable variations* if $\sum_{n=1}^{\infty} \text{var}_n(f) < \infty$, and that f *depends only on the future* if $x_i = y_i$ for all $i \geq 0$ implies that $f(x) = f(y)$.

For a suspension flow $\{S_t\} = (T, f)$ and for a set $C \subset X$, we put

$$\tilde{\tau}(x, C) = \inf\{t > 0 : S_t(x, 0) \in C \times \{0\}\}, \quad x \in X, \quad (2.3)$$

so that $\tilde{\tau}(x, C)$ is the first hitting time of $C \times \{0\}$ for a point $x \times \{0\} \in X_f$.

Theorem 2.2. *For a countable alphabet topological Markov shift (X, T) corresponding to a connected graph Γ , let $f : X \rightarrow [c, \infty)$, $c > 0$, be a function with summable variations depending only on the future, and let $\{S_t\} = (T, f)$ be the suspension flow constructed by T and f . Assume that $\bar{\mu} \in \mathcal{M}_{T,f}$ is a measure positive on all cylinders in X , and that for each $l > 0$, there exists a simple word $w \in W(\Gamma)$ with $|w| > l$ such that for every word $\hat{w}$ that does not contain w and for $\bar{\mu}$ -almost all $x \in C_{w\hat{w}w}$, we have*

$$|\bar{\mu}(C_{w\hat{w}w})/\bar{\mu}(C_w) - e^{-s\tilde{\tau}(x, C_w)}| \leq e^{-\alpha|w| - s\tilde{\tau}(x, C_w)}, \quad (2.4)$$

where $\tilde{\tau}(x, C_w)$ is defined in (2.3) and α, s are positive constants (depending only on $\bar{\mu}$). Then

- (i) $s = h_{top}(T, f)$,
- (ii) if $s = h(T, f; \bar{\mu}_f)$, then $\bar{\mu}_f$ is the unique measure of maximal entropy for the flow $\{S_t\} = (T, f)$.

Remark 2.3. The ratio in the left hand side of (2.4) is clearly the conditional measure of $C_{w\hat{w}w}$, given C_w .

Remark 2.4. The assumption that f depends only on the future is made just for convenience: Theorem 2.2 remains true without this assumption, but in the sequel we use it only in the above particular form.

Let us outline the proof of Theorem 2.2. At the first stage we consider (in Subsection 2.2) the particular case where (X, T) is a Bernoulli shift, while $f(x)$, $x \in X$, depends only on x_0 (we then say that f depends on the zeroth coordinate). In this case the topological entropy $h_{top}(T, f)$ can be expressed explicitly in terms of f . At the next stage we come back to the general case and prove that the supremum in the definition of $h_{top}(T, f)$ can be taken over ergodic measures that are positive on all cylinders in X (see Subsection 2.4). This enables us to state that $h_{top}(T, f) = h_{top}(T_C, f_C)$ for the suspension flow (T_C, f_C) where T_C is the transformation induced by T on a cylinder $C \subset X$ and f_C is determined naturally by f , T , and C . If we chose $C = \{x : (x_0, \dots, x_k) = w\}$, where $w \in W(\Gamma)$, then (C, T_C) is isomorphic to the countable alphabet Bernoulli shift. Hence the flow (T_C, f_C) is isomorphic to a suspension flow (σ, φ) built over this Bernoulli shift. Here we use a finite approximation and find a function φ^w that depends on the zeroth coordinate and is uniformly close to φ (when w is long enough). The topological entropies of the suspension flows (σ, φ) and (σ, φ^w) are also close to each other. We apply the results obtained at the first stage to the latter flow and rewrite inequality (2.4) for it (see Subsection 2.5). By directing the length of w to infinity we complete the proof of the equality $s = h_{top}(T, f)$, which implies that the measure under consideration has maximal entropy. We reduce the uniqueness of such a measure to that of the corresponding equilibrium measure and here use the uniqueness theorem by Buzzi and Sarig [6].

2.2 Entropy of suspension flows over Bernoulli shifts

The proof of Theorem 2.2 is based essentially on some properties of suspension flows constructed by a topological Markov shift (in particular, by a Bernoulli shift) and functions of one or finitely many coordinates. Some of

these properties, studied first by Savchenko [20], are described in this section. We include proofs for the reader's convenience. Our approach is close to that of [20].

We begin with two simple lemmas. Let $\mathcal{N} = \mathbb{N}$ or $\{1, \dots, n\}$, $n \geq 2$, and let $\mathbf{c} = (c_i, i \in \mathcal{N})$ be a sequence of real numbers such that $\inf_{i \in \mathcal{N}} c_i > 0$. Denote by $\mathcal{P} = \mathcal{P}_{\mathcal{N}, \mathbf{c}}$ the family of sequences $\mathbf{p} = (p_i, i \in \mathcal{N})$ such that

$$p_i \geq 0 \ (i \in \mathcal{N}), \quad \sum_{i \in \mathcal{N}} p_i = 1, \quad \sum_{i \in \mathcal{N}} p_i c_i < \infty. \quad (2.5)$$

(Certainly, $\mathcal{P}_{\mathcal{N}, \mathbf{c}}$ does not depend on $\mathbf{c}$ when $|\mathcal{N}| < \infty$.) Let

$$H(\mathbf{p}) = H_{\mathcal{N}, \mathbf{c}}(\mathbf{p}) := - \left(\sum_{i \in \mathcal{N}} p_i \log p_i \right) \left(\sum_{i \in \mathcal{N}} p_i c_i \right)^{-1}, \quad \mathbf{p} \in \mathcal{P} \quad (2.6)$$

(we as usual let $0 \log 0 = 0$).

Lemma 2.5. *If $\mathbf{p} \in \mathcal{P}$ is such that $p_k = 0$ for some $k \in \mathcal{N}$, then there exists $\mathbf{p}' = (p'_i, i \in \mathcal{N}) \in \mathcal{P}$ with $p'_i > 0$ for all i such that $H(\mathbf{p}') \geq H(\mathbf{p})$, where the inequality is strict when $H(\mathbf{p}) < \infty$.*

Proof. We divide $\mathcal{N}$ into two non-empty subsets, $\mathcal{N}^0 = \{i \in \mathcal{N} : p_i = 0\}$ and $\mathcal{N}^1 = \mathcal{N} \setminus \mathcal{N}^0$. Fix an arbitrary $l \in \mathcal{N}^1$ and for $t \in [0, p_l]$ let $\mathbf{p}^t = (p_i^t, i \in \mathcal{N})$, where $p_k^t = t$, $p_l^t = p_l - t$, and $p_i^t = p_i$ for $i \neq k, l$. (By assumption, $k \in \mathcal{N}^0$.) Clearly, $\mathbf{p}^t \in \mathcal{P}$ and $H(\mathbf{p}^t) = \infty$ when $H(\mathbf{p}) = \infty$. A simple calculation shows that if $H(\mathbf{p}) < \infty$, then the right-hand derivative $\frac{d^+}{dt} H(\mathbf{p}^t)$ at $t = 0$ is $+\infty$. Hence $H(\mathbf{p}^t) > H(\mathbf{p})$ when $t > 0$ is small enough.

If $\mathcal{N}^0 = \{k\}$, the proof is completed. If $\mathcal{N}^0 \setminus \{k\} \neq \emptyset$, we first consider the case $H(\mathbf{p}) < \infty$. Fix an arbitrary $t \in (0, p_l)$ for which $H(\mathbf{p}^t) > H(\mathbf{p})$. It is easy to find positive numbers q_i , $i \in \mathcal{N}^0 \setminus \{k\}$, such that

$$\sum_{i \in \mathcal{N}^0 \setminus \{k\}} q_i = 1, \quad \sum_{i \in \mathcal{N}^0 \setminus \{k\}} q_i (c_i - \log q_i) < \infty. \quad (2.7)$$

For $s \in [0, t]$ we put $\mathbf{p}^{t,s} = (p_i^{t,s}, i \in \mathcal{N})$, where

$$p_i^{t,s} = sq_i, \ i \in \mathcal{N}^0 \setminus \{k\}; \quad p_i^{t,s} = p_i, \ i \in \mathcal{N}^1 \setminus \{l\}; \quad p_k^{t,s} = t - s; \quad p_l^{t,s} = p_l - t. \quad (2.8)$$

From (2.7) it follows that $\mathbf{p}^{t,s} \in \mathcal{P}$ and $\lim_{s \rightarrow 0} H(\mathbf{p}^{t,s}) = H(\mathbf{p}^t)$. Therefore, $H(\mathbf{p}^{t,s}) > H(\mathbf{p})$ as $s > 0$ is small enough, and since $p_i^{t,s} > 0$, we can take $\mathbf{p}^{t,s}$ with one of these s for $\mathbf{p}'$.

It remains to note that if $H(\mathbf{p}) = \infty$, then $H(\mathbf{p}^{t,s}) = \infty$ for all $s \in [0, t)$ (see (2.7), (2.8)). $\square$

Lemma 2.6. *Let $\mathcal{N} = \{1, \dots, n\}$, $n \geq 2$, and let $\mathbf{c}$, $\mathcal{P} = \mathcal{P}_{\mathcal{N}}$, $H = H_{\mathcal{N}, \mathbf{c}}$ be as above. Then $\sup_{\mathbf{p} \in \mathcal{P}} H(\mathbf{p})$ is the unique solution to the equation $F_n(\beta) = 1$, where $F_n(\beta) = \sum_{i=1}^n e^{-\beta c_i}$.*

Proof. Since H is a continuous function on the compact set $\mathcal{P} \subset \mathbb{R}^n$, its supremum is attained at a point $\mathbf{p}^0 = (p_i^0, i = 1, \dots, n) \in \mathcal{P}$. By Lemma 2.5 $p_i^0 > 0$ for all i . Let $\mathcal{P}^+ = \{\mathbf{p} \in \mathcal{P} : p_i > 0, i = 1, \dots, n\}$. For $\mathbf{p} \in \mathcal{P}^+$ we put $p_1 = 1 - \sum_{i=2}^n p_i$ and consider the equations $\partial H(\mathbf{p}) / \partial p_i = 0$, $i \in \mathcal{N} \setminus \{1\}$. From this system we derive that if $\mathbf{p}^0$ is a point of extremum of $H(\mathbf{p})$, then $p_i^0 = e^{-\beta^0 c_i} / F_n(\beta^0)$, $1 \leq i \leq n$, where $\beta^0 = \text{const} > 0$. Hence the statement we are proving is true when $c_i = c_1$ for $i = 2, \dots, n$. Otherwise we take any i for which $c_i \neq c_1$ and from the equation $\frac{\partial H(\mathbf{p})}{\partial p_i} \big|_{\mathbf{p}=\mathbf{p}^0} = 0$, where $p_1 = 1 - \sum_{i=2}^n p_i$, obtain $\beta^0 = H(\mathbf{p}^0)$. On the other hand, by substituting $\mathbf{p}^0$ for $\mathbf{p}$ in $H(\mathbf{p})$ we see that $H(\mathbf{p}^0) = \beta^0 - \frac{F_n(\beta^0)}{F_n'(\beta^0)} \log F_n(\beta^0)$. Therefore, $\log F_n(\beta^0) = 0$, i.e., β^0 is a root of the equation $F_n(\beta) = 1$. This root is unique, since $F_n(\beta)$ decreases in β . Finally, $H(\mathbf{p}^0) = \max_{\mathbf{p} \in \mathcal{P}} H(\mathbf{p})$, because, as was mentioned above, every point of maximum belongs to $\mathcal{P}^+$, hence the equations $\partial H(\mathbf{p}) / \partial p_i = 0$, $i \in \mathcal{N} \setminus \{1\}$, $p_1 = 1 - \sum_{i=2}^n p_i$ must hold at this point. But we already know that these equations have only one solution. $\square$

Let us now consider a countable alphabet topological Bernoulli shift (X, T) with $X = V^{\mathbb{Z}}$, and the suspension flow $\{S_t\} = (T, f)$ constructed by T and a function f such that $f(x) = f_0(x_0)$, $x = (x_i, i \in \mathbb{Z})$, where $f_0 : V \rightarrow [c, \infty)$, $c > 0$. Let

$$F(\beta) = \sum_{v \in V} e^{-\beta f_0(v)}, \quad \beta \geq 0.$$

Lemma 2.7. *If there exists $\beta_0 \geq 0$ with $F(\beta_0) = 1$, then $h_{\text{top}}(T, f) = \beta_0$. Otherwise $h_{\text{top}}(T, f) = \sup\{\beta \geq 0 : F(\beta) = \infty\}$.*

Proof. Denote by $B_{T,f}$ the family of all Bernoulli measures in $\mathcal{M}_{T,f}$. Each $\nu \in B_{T,f}$ is determined by the one-dimensional distribution $\{p^\nu(v), v \in V\}$,

where

$$p^\nu(v) = \nu(C_v) \geq 0, \quad \sum_{v \in V} p^\nu(v) = 1, \quad \sum_{v \in V} p^\nu(v) f_0(v) < \infty.$$

We note that

$$\sup_{\mu \in \mathcal{M}_{T,f}} [h(T; \mu)/\mu(f)] = \sup_{\nu \in B_{T,f}} [h(T; \mu)/\mu(f)]. \quad (2.9)$$

Indeed, every $\mu \in \mathcal{M}_{T,f}$ gives rise to the measure $\mu_B \in B_{T,f}$ with $p^{\mu_B}(v) = \mu_B(C_v) = \mu(C_v)$. Clearly, $\mu_B(f) = \mu(f)$, and basic properties of the measure-theoretic entropy imply that $h(T; \mu) \leq h(T; \mu_B)$.

Let us number in an arbitrary way the elements $v \in V$ and put $B^{(n)} = \{\nu \in B_{T,f} : p^\nu(v_i) = 0 \text{ for } i \geq n+1\}$, $n \in \mathbb{N}$. For each $\mu \in B_{T,f}$, one can easily find a sequence of measures $\nu_n \in B^{(n)}$ such that

$$\lim_{n \rightarrow \infty} [h(T; \nu_n)/\nu_n(f)] = h(T; \mu)/\mu(f).$$

Therefore, by (2.9),

$$\sup_{\mu \in \mathcal{M}_{T,f}} [h(T; \mu)/\mu(f)] = \sup_{n \in \mathbb{N}} \sup_{\nu \in B^{(n)}} [h(T; \nu)/\nu(f)]. \quad (2.10)$$

We now notice that the relations

$$p_i := p^\nu(v_i), \quad 1 \leq i \leq n; \quad \mathbf{p} = \mathbf{p}^\nu := (p_1, \dots, p_n)$$

establish a one-to-one correspondence between $B^{(n)}$ and $\mathcal{P} = \mathcal{P}_{\mathcal{N}}$ with $\mathcal{N} = \{1, \dots, n\}$, and that $h(T; \nu)/\nu(f) = H_{\mathcal{N}, \mathbf{c}}(\mathbf{p}) = H(\mathbf{p})$, where $\mathbf{c} = (c_i, i \in \mathcal{N})$, $c_i = f_0(v_i)$, $i \in \mathcal{N}$ (see (2.5) and (2.6)).

By Lemma 2.6 the right-hand side of (2.10) is $\sup_n \beta_n$, where β_n is determined by $F_n(\beta_n) = 1$. Let us note that F_n is the n th partial sum of the series for F and that both F_n and F are strictly decreasing functions (for F it is true on the semi-axis where F is finite). Hence $\sup_n \beta_n = \lim_{n \rightarrow \infty} \beta_n$. We consider two possible cases and first suppose that $F(\beta) = \infty$ for all $\beta \geq 0$. It is clear that in this case $\lim_{n \rightarrow \infty} \beta_n = \infty$. Otherwise there exists a unique $\beta_\infty > 0$ such that either $F(\beta_\infty) = 1$, or $F(\beta) < 1$ for $\beta \geq \beta_\infty$ and $F(\beta) = \infty$ for $\beta < \beta_\infty$. Since $F_n(\beta) < F_{n+1}(\beta) < F(\beta)$ for all $n \geq 1$ and $\beta \geq 0$, in both cases we have $\lim_{n \rightarrow \infty} \beta_n \leq \beta_\infty$. If $\lim_{n \rightarrow \infty} \beta_n =: \beta'_\infty < \beta_\infty$, then $F(\beta'_\infty) > 1$ (in the latter case $F(\beta'_\infty) = \infty$). Therefore $F_n(\beta'_\infty) > 1$ for n large enough. But $\beta'_\infty > \beta_n$, hence $F_n(\beta'_\infty) < F_n(\beta_n) < 1$ for all n . From this we conclude that $\lim_{n \rightarrow \infty} \beta_n = \beta_\infty$. We thus come to both statements of the lemma. $\square$

2.3 Induced automorphisms and Markov-Bernoulli reduction

For the next lemma we have to remind the following definition. Let T be an automorphism of the space $(X, \mathcal{B})$, and let $C \in \mathcal{B}$. Denote

$$X_C = \{x \in X : \sum_{n < 0} \mathbf{1}_C(T^n x) = \sum_{n > 0} \mathbf{1}_C(T^n x) = \infty\}, \quad C' = C \cap X_C. \quad (2.11)$$

Thus C' consists of all points in C that return to C infinitely often in forward and backward time. Let also

$$\tau(T, C; x) = \min\{n > 0 : T^n x \in C\}, \quad T_{C'} x = T^{\tau(T, C; x)} x, \quad x \in C'. \quad (2.12)$$

It is clear that the sets X_C, C' are measurable and invariant with respect to T and $T_{C'}$ respectively, and that $T_{C'}$ is an automorphism of the set C' provided with the induced Borel σ -algebra; $T_{C'}$ is said to be the *induced automorphism* on C' .

Lemma 2.8. *Let (T, f) be the suspension flow constructed by an automorphism T of $(X, \mathcal{B})$ and a $\mathcal{B}$ -measurable function $f : X \rightarrow [c, \infty)$, $c > 0$, and let $C \in \mathcal{B}$. Then the suspension flow $(T|_{X_C}, f|_{X_C})$ constructed by the restrictions of T and f to X_C is isomorphic to the suspension flow $(T_{C'}, f_{C'})$, where*

$$f_{C'}(x) = \sum_{i=0}^{\tau(T, C; x)-1} f(T^i x), \quad x \in C'. \quad (2.13)$$

Furthermore, if $\mu \in \mathcal{M}_{T, f}$ is ergodic and such that $\mu(C) > 0$, then $\mu(f) = \int_{C'} f_{C'} d\mu$ and the suspension flow $(T, f; \mu_f)$ is isomorphic to the suspension flow $(T_{C'}, f_{C'}; (\mu_{C'})_{f_{C'}})$, where μ_f is the f -lifting of μ , $\mu_{C'}$ is the normalized restriction of μ to C' , and $(\mu_{C'})_{f_{C'}}$ is the $f_{C'}$ -lifting of $\mu_{C'}$.

We omit the proof of this lemma, since it follows immediately from standard facts of ergodic theory (see, for instance, [7]).

The following construction is reminiscent of the Doeblin first return method in the theory of Markov chains and has appeared repeatedly in the literature in different forms (presumably for the first time — in [9], see also [10] and [12]).

Let $w = (v_1, \dots, v_l) \in W(\Gamma)$ and $C = C_w$. Then X_C defined by (2.11) can be described as follows: $x \in X$ belongs to X_C if and only if there is an

increasing sequence of integers $i_k = i_k(x)$, $-\infty < k < \infty$, such that $i_k \leq 0$ for $k \leq 0$, $i_k > 0$ for $k > 0$, and $(x_{i_k}, \dots, x_{i_k+l-1}) = w$ for every k , while no other segment of x agrees with w . Furthermore, C' consists of those x for which $i_0(x) = 0$. It is clear that

$$i_1(x) = \tau(T, C'; x), \quad i_k(x) \geq i_1(x) + k - 1, \quad x \in C'.$$

Denote by A_w the set of all words $w' = (v'_1, \dots, v'_{l'}) \in W(\Gamma)$ with $l' > l$ such that $(v'_1, \dots, v'_l) = (v'_{l'-l+1}, \dots, v'_{l'}) = w$ and no other subword of w' (i.e., a word of the form $(v'_m, v'_{m+1}, \dots, v'_n)$, $1 \leq m \leq n \leq l'$) agrees with w . It is easy to see that if $x \in X_C$, then for each $k \in \mathbb{Z}$, the word $(x_{i_k}, x_{i_k+1}, \dots, x_{i_k+l-1})$ belongs to A_w . We thus obtain a mapping $\Psi_w : X_C \rightarrow (A_w)^\mathbb{Z}$ measurable with respect to the appropriate Borel σ -algebras; its restriction to C' obviously induces a one-to-one correspondence between C' and $(A_w)^\mathbb{Z}$. Moreover, if $x \in C'$, then $\Psi_w T_{C'} x = \sigma_w \Psi_w x$, where σ_w is the shift transformation on $Y_w := (A_w)^\mathbb{Z}$, i.e., $(\sigma_w y)_i = y_{i+1}$, $y = (y_i, i \in \mathbb{Z}) \in Y_w$. Therefore, $T_{C'}$ is isomorphic to the countable alphabet Bernoulli shift (Y_w, σ_w) with alphabet A_w . Here and in the sequel we consider each $a \in A_w$ as either a word in the alphabet V or a letter in the new alphabet A_w . What of these two possibilities takes place will always be clear from the context.

This construction reduces in essence the study of the topological Markov shift (X, T) to that of a topological Bernoulli shift determined by w , and so we shall refer to it as the *Markov–Bernoulli (M–B) reduction* applied to (X, T) and w .

2.4 Positive measures

Our next aim is to show that the topological entropy of a suspension flow over a Markov shift can be computed using only ergodic measures that are positive on all cylinders.

Lemma 2.9. *Let (X, T) and f be as in Theorem 2.2 and let (T, f) be the suspension flow constructed by T and f . Then*

$$h_{\text{top}}(T, f) = \sup_{\mu \in \mathcal{E}_{T,f}^+} h(T, f; \mu_f),$$

where $\mathcal{E}_{T,f}^+$ consists of all ergodic measures in $\mathcal{M}_{T,f}$ that are positive on all cylinders in X .

Proof. Denote by $\mathcal{E}_{T,f}$ the set of ergodic measures in $\mathcal{M}_{T,f}$. If $\mu \in \mathcal{M}_{T,f} \setminus \mathcal{E}_{T,f}$, i.e., if μ is non-ergodic with respect to T , then μ_f , the f -lifting of μ , is non-ergodic with respect to the suspension flow (T, f) . The flow $(T, f; \mu_f)$ can be decomposed into ergodic components (see [17]). This means the following. There exists a measurable partition ζ of the space (X_f, μ_f) such that μ_f -almost every element C_ζ of ζ is (T, f) -invariant and the conditional measure $(\mu_f)^{C_\zeta}$ induced by μ on C_ζ is invariant and ergodic with respect to the restriction of (T, f) to C_ζ . We may consider $(\mu_f)^{C_\zeta}$ as a measure on the whole space X_f ; it is (T, f) -invariant and ergodic. By a general formula ([18], Section 9)

$$h(T, f; \mu_f) = \int_{X_f|\zeta} h(T, f; (\mu_f)^{C_\zeta}) \mu_{f,\zeta}(dC_\zeta),$$

where $\mu_{f,\zeta}$ is the projection of μ_f on the quotient space $X_f|\zeta$. Hence, for every $\varepsilon > 0$, there is an element C_ζ with $h(T, f; (\mu_f)^{C_\zeta}) > h(T, f; \mu_f) - \varepsilon$. On the other hand, $(\mu_f)^{C_\zeta}$, being a (T, f) -invariant probability measure on X_f , is the f -lifting of a T -invariant probability measure μ^{C_ζ} on X , i.e., $(\mu_f)^{C_\zeta} = (\mu^{C_\zeta})_f$. It is clear that $\mu^{C_\zeta}(f) < \infty$ and (T, μ^{C_ζ}) is ergodic. Since $\varepsilon > 0$ was arbitrary small, we conclude that

$$h_{\text{top}}(T, f) = \sup_{\mu \in \mathcal{E}_{T,f}} h(T, f; \mu_f).$$

Let $\mathcal{E}_{T,f}^0 := \mathcal{E}_{T,f} \setminus \mathcal{E}_{T,f}^+$ and assume that, contrary to the lemma we have to prove, for some $\delta \in (0, \infty)$,

$$\sup_{\mu \in \mathcal{E}_{T,f}^0} h(T; \mu)/\mu(f) > \sup_{\mu \in \mathcal{E}_{T,f}^+} h(T; \mu)/\mu(f) + \delta, \quad (2.14)$$

which in particular means that

$$\sup_{\mu \in \mathcal{E}_{T,f}^+} h(T; \mu)/\mu(f) < \infty.$$

By virtue of (2.14) there is $\mu^0 \in \mathcal{E}_{T,f}^0$ such that

$$h(T; \mu^0)/\mu^0(f) \geq \sup_{\mu \in \mathcal{E}_{T,f}^+} h(T; \mu)/\mu(f) + \delta/2. \quad (2.15)$$

To show that this is impossible we first consider the case $h(T; \mu^0) < \infty$ and let $h^0 = h(T, \mu^0)/\mu^0(f)$. Since f has summable variations, one can find

$n_\delta \in \mathbb{N}$ such that, for every $n \geq n_\delta$, there is a function $f_n : X \rightarrow \mathbb{R}_+$ with the following three properties: $f_n(x) = f_n(y)$ whenever $x_i = y_i$ for $|i| \leq n$, $\inf_{x \in X} f_n(x) \geq c$, and

$$\sup_{x \in X} |f(x) - f_n(x)| \leq \delta c^2 / 8h^0.$$

One can easily check that then

$$\left| \frac{h(T; \mu)}{\mu(f_n)} - \frac{h(T; \mu)}{\mu(f)} \right| < \delta \quad (2.16)$$

for every $\mu \in \mathcal{E}_{T,f}^+ \cup \{\mu^0\}$. Hence (see (2.15))

$$h(T; \mu^0) / \mu^0(f_n) \geq \sup_{\mu \in \mathcal{E}_{T,f_n}^+} h(T; \mu) / \mu(f_n) + \delta/4. \quad (2.17)$$

Since $|f - f_n| < \text{const}$, the functions f and f_n are integrable or not integrable with respect to a finite measure simultaneously. Hence $\mathcal{E}_{T,f_n} = \mathcal{E}_{T,f}$ and $\mathcal{E}_{T,f_n}^+ = \mathcal{E}_{T,f}^+$.

If $h(T; \mu^0) = \infty$, then (2.17) clearly holds as well.

Using the assumption $\mu^0 \in \mathcal{E}_{T,f}^0$, we find a word $w^0 \in W(\Gamma)$ with $\mu^0(C_{w^0}) = 0$. Fix an arbitrary $n^1 \geq \max\{n_\delta, |w^0|\}$ and a word $w^1 \in W(\Gamma)$ with $|w^1| = n^1$, $\mu^0(w^1) > 0$. Then we set $f^1 := f_{n^1}$, $C := C_{w^1}$, and apply the M-B reduction to (X, T) and w^1 . By Lemma 2.8 the suspension flow $(T|_{X_C}, f^1|_{X_C})$ is isomorphic to the suspension flow $(\sigma, \varphi) := (\sigma_{w^1}, \varphi_{f^1, w^1})$, where

$$\varphi(y) := (f^1)_{C'}(\Psi_{w^1}^{-1}y), \quad y \in Y_{w^1}. \quad (2.18)$$

Notice that the function φ is constant on every one-dimensional cylinder $\{y \in Y : y_0 = a\}$, $a \in A_{w^1}$; the reason is that each $a \in A_{w^1}$ when considered as a word from $W(\Gamma)$ is not shorter than w^1 .

Let us carry over the measure $\mu_{C'}^0$, the normalized restriction of μ^0 to C' (where C' is defined in (2.11)), to Y via the mapping Ψ_{w^1} to obtain a Borel probability measure ν^0 on Y . From the above-described properties of Ψ_{w^1} it follows that the automorphisms $(T_{C'}; (\mu^0)_{C'})$ and $(\sigma; \nu^0)$ are isomorphic and hence, by Lemma 2.8, the suspension flow $(T, f^1; (\mu^0)_{f^1})$ is isomorphic to the suspension flow $(\sigma, \varphi; (\nu^0)_\varphi)$, where $(\mu^0)_{f^1}$ and $(\nu^0)_\varphi$ are the f^1 -lifting of μ^0 and the φ -lifting of ν^0 , respectively. Therefore, by (2.1),

$$h(T; \mu^0) / \mu^0(f^1) = h(\sigma; \nu^0) / \nu^0(\varphi). \quad (2.19)$$

If we change ν^0 for a σ -invariant Bernoulli measure ν^1 with the same one-dimensional distribution (i.e., with $\nu^1(C_a) = \nu^0(C_a)$ for all $a \in A_{w^1}$, where $C_a = \{y \in Y : y_0 = a\}$), then the numerator on the right-hand side of (2.19) can only increase, while the denominator will not change (since φ is constant on every cylinder C_a , $a \in A_{w^1}$).

From the definition of ν^0 and ν^1 it follows that $\nu^0(C_{a^0}) = \nu^1(C_{a^0}) = 0$ for some $a^0 \in A_{w^1}$. Indeed, let $w^1 = (v_1^1, \dots, v_{l_1}^1)$. Since the graph Γ is connected, there exists a word $(v_1, \dots, v_r) \in W(\Gamma)$ with $(v_1, \dots, v_{l_1}) = w^1$, $(v_{r-l_0+1}, \dots, v_r) = w^0$, where $l_0 = |w^0|$. Choose an arbitrary shortest word of this type and denote it by w' . Similarly, let w'' be one of the shortest words in which there are an initial subword and a terminal subword that coincide with w^0 and w^1 , respectively. From the assumption that $\mu^0(w^0) = 0$, $\mu^0(w^1) > 0$, $|w^0| \leq |w^1|$ it follows that $w'' = w^0 \hat{w}$ where $\hat{w}$ can be of one of the following three forms: (a) $\hat{w} = w^1$; (b) $\hat{w} = \hat{w}^1 w^1$, $\hat{w}^1 \in W(\Gamma)$; (c) $\hat{w} = (v_k^1, \dots, v_{l_1}^1)$, $1 < k \leq l_1$, is a terminal subword of w^1 . Consider the word $w' \hat{w}$. One easily checks that $w' \hat{w} \in A_{w^1}$. Moreover, $\mu^0(w' \hat{w}) = 0$, because $w' \hat{w}$ contains w^0 as a subword. Hence $\mu_{C'_{w^1}}^0(C'_{w^1} \cap C_{w' \hat{w}}) = 0$. We can put $a^0 := w' \hat{w}$. Since $\Psi_{w^1}(C' \cap C_{w' \hat{w}}) = C_{a^0}$, we have $\nu^1(C_{a^0}) = \nu^0(C_{a^0}) = 0$.

We now want to perturb ν^1 within the class of Bernoulli measures on Y in such a way as to obtain a measure for which the right-hand side of (2.19) is bigger than for ν^0 and which is positive on all cylinders.

Since $\varphi(y)$, $y = (y_i, i \in \mathbb{Z}) \in Y$, depends solely on y_0 , we have $\varphi(y) = \varphi_0(y_0)$, where φ_0 is a function on A_{w^1} .

Using Lemma 2.5, we find a σ -invariant Bernoulli measure ν^2 on Y such that if $h(\sigma; \nu^1) < \infty$, then

$$\frac{h(\sigma; \nu^2)}{\nu^2(\varphi)} > \frac{h(\sigma; \nu^1)}{\nu^1(\varphi)} \geq \frac{h(\sigma; \nu^0)}{\nu^0(\varphi)}, \quad (2.20)$$

and if $h(\sigma; \nu^1) = \infty$, then $h(\sigma; \nu^2) = \infty$ as well.

Apply the mapping $\Psi_{w^1}^{-1}$ to transfer the measure ν^2 to C' and denote the resulting measure by μ' . The suspension flow $(\sigma, \varphi; (\nu^2)_\varphi)$ is then isomorphic to the suspension flow $(T_{C'}, (f^1)_{C'}; (\mu')_{(f^1)_{C'}})$. Let $C'(n) = \{x \in C' : \tau_{T, C'}(x)\}$, $n = 1, 2, \dots$, and $\mu'_n = \mu'|_{C'(n)}$ be the restriction of μ' to $C'(n)$ considered as a measure on X . Then the measure

$$\mu'' := \sum_{n=1}^{\infty} \sum_{k=0}^{n-1} T^k \circ \mu'_n$$

is concentrated on X_C and T -invariant. By normalizing μ'' we obtain a probability measure μ''' . By Lemma 2.8 the flows $(T|_{X_C}; f^1|_{X_C})$ and $(T_{C'}, (f^1)_{C'})$ are isomorphic. Then the flow $(T, f_1; (\mu''')_{f^1})$ is isomorphic to the flow $(T_{C'}, (f^1)_{C'}; (\mu')_{f^1|_{C'}})$ and hence (see above) to the flow $(\sigma, \varphi; (\nu^2)_\varphi)$. Therefore,

$$h(T; \mu''')/\mu'''(f^1) = h(\sigma; \nu^2)/\nu^2(\varphi) > h(T; \mu^0)/\mu^0(f^1) \quad (2.21)$$

(see (2.19), (2.20)). It is clear that $\mu''' \in \mathcal{E}_{T,f}$. Moreover, $\mu''' \in \mathcal{E}_{T,f}^+$. Otherwise we could apply to μ''' the procedure that lead us to the measure ν^0 , starting from μ^0 . The resulting measure would coincide with $\nu^{t,s}$, and there would be a letter $a \in A_{w^1}$ with $\nu^{t,s}(C_a) = 0$. But we know that this is impossible. Thus (2.21) contradicts (2.17) with $f_n = f^1$ and hence contradicts (2.14). $\square$

Corollary 2.10. *Let Γ , (X, T) , f be as in Theorem 2.2, (Y_w, σ_w) be the topological Bernoulli shift obtained from (X, T) and w by the M - B reduction, where $w \in W(\Gamma)$, and let $\varphi_{f,w}$ be the function defined in (2.18). Then the suspension flows (T, f) and $(\sigma_w, \varphi_{f,w})$ have the same topological entropy.*

Proof. As before, we let $C = C_w$ and use the notation in (2.11)–(2.13). From the definition of σ_w and $\varphi_{f,w}$ it follows immediately that the suspension flows $(T_{C'}, f_{C'})$ and $(\sigma_w, \varphi_{f,w})$ are isomorphic and hence $h_{\text{top}}(T_{C'}, f_{C'}) = h_{\text{top}}(\sigma_w, \varphi_{f,w})$. Similarly, by virtue of Lemma 2.8, $h_{\text{top}}(T_{C'}, f_{C'}) = h_{\text{top}}(T|_{X_C}, f|_{X_C})$. But $h_{\text{top}}(T|_{X_C}, f|_{X_C}) \leq h_{\text{top}}(T, f)$, because X_C is a T -invariant subset of X . Hence

$$h_{\text{top}}(\sigma_w, \varphi_{f,w}) = h_{\text{top}}(T_{C'}, f_{C'}) \leq h_{\text{top}}(T, f). \quad (2.22)$$

On the other hand, by the same Lemma 2.8 $h(T_{C'}, f_{C'}; \tilde{\mu}_{C'}) = h(T, f; \tilde{\mu})$ for every $\mu \in \mathcal{E}_{T,f}^+$, where $\tilde{\mu}_{C'}$ is the $f_{C'}$ -lifting of the normalized restriction of μ to C' , and $\tilde{\mu}$ is the f -lifting of μ . The supremum in $\mu \in \mathcal{E}_{T,f}^+$ of the left-hand side of the last equality is clearly not bigger than $h(T_{C'}, f_{C'})$, while by Lemma 2.9 the supremum of the right-hand side is $h_{\text{top}}(T, f)$. Hence $h_{\text{top}}(T_{C'}, f_{C'}) \geq h_{\text{top}}(T, f)$, which together with (2.22) yields what we are proving. $\square$

2.5 Proof of Theorem 2.2

Fix notation as in Theorem 2.2. Let us also fix n and for a while write w and C instead of w_n and C_{w_n} , respectively. Consider the sets X_C , C' , the

induced transformation $T_{C'} : C' \rightarrow C'$, and the function $f_{C'}$ (see (2.11)–(2.13)). Apply the M–B reduction to (X, T) and w .

From (2.3), (2.13) it is clear that $\tilde{\tau}(T, C; x) = f_{C'}(x)$ for every $x \in C'$. Thus (2.4) can be rewritten in the form

$$|\bar{\mu}(C_{w\hat{w}w})/\bar{\mu}(C) - e^{-sf_{C'}(x)}| \leq e^{-\alpha|w|-sf_{C'}(x)}, \quad (2.23)$$

which is true for $\bar{\mu}$ -almost all $x \in C_{w\hat{w}w}$.

The simplicity of w implies that each word $a \in A_w$ is of the form $a = w\hat{w}w$, where $\hat{w} \in W(\Gamma)$ ($\hat{w}$ may be an empty word if $ww \in W(\Gamma)$), and $\hat{w}$ does not contain w as a subword.

By assumption, the measure $\bar{\mu}$ is positive on all cylinders and T -invariant. Hence $\bar{\mu}(C) = \bar{\mu}(C') > 0$, and we can normalize $\bar{\mu}$ on C' to obtain a $T_{C'}$ -invariant probability measure $\bar{\mu}'$. Its image $\nu' := (\Psi_w)_*\bar{\mu}'_0$ is a probability measure ν' on $Y = Y_w$ invariant with respect the shift transformation σ . From the definition of Ψ_w it follows that, for $a := w\hat{w}w \in A_w$,

$$C_a := \{y \in Y : y_0 = a\} = \Psi_w C_{w\hat{w}w}$$

and hence

$$\nu'(C_a) = \mu_0(C_{w\hat{w}w})/mu_0(C_w). \quad (2.24)$$

Notice that $\nu'(C_a) > 0$ for all $a \in A_w$.

Taking into account the relation between $f_{C'}$ and $\varphi = \varphi_{f,w}$ (see (2.18)) and using (2.23), (2.24), we obtain

$$|\nu'(C_a) - e^{-s\varphi(y)}| \leq e^{-\alpha|w|-s\varphi(y)}, \quad a \in A_w, \quad (2.25)$$

for ν' -almost all $y \in C_a$.

Our next step is to approximate φ by a function that is constant on each cylinder C_a , $a \in A_w$.

Since w is simple, we have $\tau(T, C'; x) \geq |w|$ for all $x \in C'$ (see (2.12)). We say that $x^{(1)} = (x_i^{(1)}, i \in \mathbb{Z}) \in C'$ and $x^{(2)} = (x_i^{(2)}, i \in \mathbb{Z}) \in C'$ are equivalent ($x^{(1)} \sim x^{(2)}$) if $\tau(T, C'; x^{(1)}) = \tau(T, C'; x^{(2)})$ and $x_i^{(1)} = x_i^{(2)}$ for $0 \leq i \leq \tau(T, C'; x^{(1)})$. If $x^{(1)} \sim x^{(2)}$, then (because w is simple) $x_i^{(1)} = x_i^{(2)}$ for $\tau(T, C'; x^{(1)}) \leq i \leq \tau(T, C'; x^{(1)}) + |w| - 1$ as well, from which we obtain (see (2.13))

$$\begin{aligned} & |f_{C'}(x^{(1)}) - f_{C'}(x^{(2)})| \leq \\ & \leq \sum_{i=0}^{\tau(T, C'; x)-1} |f(T^i x^{(1)}) - f(T^i x^{(2)})| \leq \sum_{n=|w|}^{\infty} \text{var}_n(f). \end{aligned} \quad (2.26)$$

Let

$$C^w(x) := \{x' \in C' : x' \sim x\}, \quad f^w(x) = \inf_{x' \in C^w(x)} f_{C'}(x'). \quad (2.27)$$

It is easy to see that $C^w(x)$ is a cylinder and that these cylinders constitute a partition of C' . Moreover, by virtue of (2.27), (2.26) the function f^w is constant on each element of this partition and

$$0 \leq f_{C'}(x) - f^w(x) \leq \sum_{n=|w|}^{\infty} \text{var}_n(f), \quad x \in C'.$$

Therefore,

$$0 \leq \varphi(y) - \varphi^w(y) \leq \sum_{n=|w|}^{\infty} \text{var}_n(f), \quad y \in Y, \quad (2.28)$$

where $\varphi^w(y) := f^w(\Psi_w^{-1}y)$ is constant on each cylinder $C_a \subset Y$, $a \in A_w$ (here, by $\Psi_w^{-1}y$ we mean the unique point $x \in C'$ such that $\Psi_w x = y$) and hence there is a function φ_0^w on A_w such that $\varphi^w(y) = \varphi_0^w(y_0)$.

With Lemma 2.7 in mind we will estimate the sum $\sum_{a \in A_w} \exp[-s\varphi_0^w(a)]$. Let

$$\delta_w := \sum_{n=|w|}^{\infty} \text{var}_n(f). \quad (2.29)$$

Since $\nu'(C_a) > 0$ for all $a \in A_w$, one can choose, for every a , a point $y_a \in C_a$ such that (2.25) holds for $y = y_a$. Hence

$$\nu'(C_a) - \exp[-\alpha|w| - s\varphi(y_a)] \leq e^{-s\varphi(y_a)} \leq \nu'(C_a) + \exp[-\alpha|w| - s\varphi(y_a)],$$

so that

$$\nu'(C_a)/(1 + e^{-\alpha|w|}) \leq e^{-s\varphi(y_a)} \leq \nu'(C_a)/(1 - e^{-\alpha|w|}), \quad a \in A_w,$$

$$1/(1 + e^{-\alpha|w|}) \leq \sum_{a \in A_w} e^{-s\varphi(y_a)} \leq 1/(1 - e^{-\alpha|w|}).$$

From (2.28), (2.29) we obtain

$$\begin{aligned}
\frac{1}{1 + e^{-\alpha|w|}} &\leq \sum_{a \in A_w} e^{-s\varphi(y_a)} \leq \sum_{a \in A_w} e^{-s\varphi_0^w(a)} \\
&= \sum_{a \in A_w} e^{-s\varphi^w(y_a)} \leq \frac{1}{1 - e^{-\alpha|w|}} + \sum_{a \in A_w} [e^{-s\varphi^w(y_a)} - e^{-s\varphi(y_a)}] \\
&= \frac{1}{1 - e^{-\alpha|w|}} + \sum_{a \in A_w} e^{-s\varphi(y_a)} [e^{s(\varphi(y_a) - \varphi^w(y_a))} - 1] \leq \frac{e^{s\delta_w}}{1 - e^{-\alpha|w|}}. \tag{2.30}
\end{aligned}$$

By assumption, we now can take $w = w_n$, where $|w_n| \rightarrow \infty$ as $n \rightarrow \infty$. From (2.30) it follows that

$$\lim_{n \rightarrow \infty} \sum_{a \in A_{w_n}} \exp[-s\varphi_0^{w_n}(a)] = 1. \tag{2.31}$$

Let

$$F_n(u) := \sum_{a \in A_{w_n}} \exp[-u\varphi_0^{w_n}(a)], \quad n = 1, 2, \dots$$

If, for a fixed n , there is a $u \in \mathbb{R}$ such that $F_n(u) = 1$ (such u can be only one), then we denote this u by u_n . Otherwise we put $u_n := \sup\{u : F_n(u) = \infty\}$. Notice that $u_n \geq 0$ (because $F_n(0) = \infty$) and $u_n < \infty$ (because of (2.30)). From the definition of φ^{w_n} it follows that $\inf_{y \in Y} \varphi^{w_n}(y) \rightarrow \infty$ as $n \rightarrow \infty$ (remind that $Y = Y_{w_n}$). Therefore, for every $\gamma > 0$, we have $dF_n(u)/du \rightarrow -\infty$ as $n \rightarrow \infty$ uniformly in u on the set $D_\gamma := \{u : \gamma < F_n(u) < \infty\}$ (we mean the right-side derivative if u is the left endpoint of the interval D_γ). Using this fact, it is easy to deduce from (2.31) that $u_n \rightarrow s$ as $n \rightarrow \infty$ (it would be sufficient to know that $dF_n(u)/du < \text{const} < 0$ on D_γ).

Let us now consider two isomorphic suspension flows, (σ, φ^{w_n}) and $(T_{C'}, f^{w_n})$. By Lemma 2.7

$$t_n = h_{\text{top}}(\sigma, \varphi^{w_n}) = h_{\text{top}}(T_{C'}, f^{w_n}), \quad n = 1, 2, \dots,$$

where $C' = (C_{w_n})'$, and hence

$$s = \lim_{n \rightarrow \infty} h_{\text{top}}(T_{C'}, f^{w_n}).$$

From (2.28) and the evident bounds $h_{\text{top}}(\sigma, \varphi^{w_n}) \leq 2s$ (as n is large enough), $\inf \varphi \geq c$, and $\inf \varphi^{w_n} \geq c$ we obtain

$$|h_{\text{top}}(\sigma, \varphi^{w_n}) - h_{\text{top}}(\sigma, \varphi)| \leq 2s\delta_n/c,$$

where $\delta_n = \sum_{k=|w_n|} \text{var}_n(f)$ (cf. (2.16)). Therefore $s = h_{\text{top}}(\sigma, \varphi)$, and hence (see Corollary 2.10) $s = h_{\text{top}}(T, f)$. So statement (i) is proved.

To prove statement (ii) suppose that $s = h_{\text{top}}(T, f; (\bar{\mu})_f)$. Together with (i) this means that $(\bar{\mu})_f$ is a measure with maximal entropy for the suspension flow (T, f) . We conclude that (T, f) can have only one measure with maximal entropy, using a result by Buzzi and Sarig [6], as follows.

Let $(\bar{\mu})_f$ be such a measure. Then by (2.2)

$$\frac{h(T, \mu)}{\mu(f)} \leq \frac{h(T, \bar{\mu})}{\bar{\mu}(f)} = s, \quad \mu \in \mathcal{M}_{T,f},$$

where $s = h_{\text{top}}(T, f)$. Hence for every $\mu \in \mathcal{M}_{T,f}$, we have $h(T, \mu) + \mu(g) \leq 0$, where $g(x) := -sf(x)$, $x \in X$, while $h(T, \bar{\mu}) + \bar{\mu}(g) = 0$, so that the topological pressure of g is zero and $\bar{\mu}$ is a g -equilibrium measure. Using the natural projection $\pi : V^{\mathbb{Z}} \rightarrow V^{\mathbb{Z}_+}$ we let $X_+ = \pi X$ and $f_+(x_+) = f(x)$ for $x_+ \in X_+$ and any $x \in \pi^{-1}x_+$ (by assumption, f is constant on the set $\pi^{-1}x_+$, so that $f(x)$ depends only on x_+). It is easily checked that $\pi T x = T_+ \pi x$, $x \in X$, where T_+ is the shift transformation on X_+ , and, moreover, that π induces a one-to-one correspondence between $\mathcal{M}_{T,f}$ and $\mathcal{M}_{T_+,f_+}$, the set of T_+ -invariant probability measures μ_+ on X_+ with $\mu_+(f_+) < \infty$. Let $\mu_{0+} \in \mathcal{M}_{T_+,f_+}$ correspond to μ_0 . Then μ_{0+} is a g_+ -equilibrium measure, where $g_+ = -sf_+$. Notice that the one-sided Markov shift T_+ is topologically transitive (because the graph Γ is connected), the topological pressure of g_+ is zero (because this is the case for g), and $\sup_{x_+ \in X_+} g_+(x_+) < 0$ (because $\inf_{x \in X} f(x) > 0$). Thus, by Theorem 1.1 from [6], there can be only one g_+ -equilibrium measure. So the proof of Theorem 2.2 is completed.

3 A covering flow

At this point we start deducing Theorem 1.1 from Theorem 2.2. The aim of the present section is to recall the construction of a flow that can be viewed (see Section 5 below) as "covering" for the Techmüller flow $\{g_t\}$. We shall show (see corollary 5.2) that our problem can be reduced to a similar problem for this covering flow (denoted by $\{P^t\}$).

We first remind some constructions due to Rauzy [16], Veech [21], and Zorich [26] (see also [24]). Using these constructions we obtain (in Subsection 3.2) the covering flow as a suspension flow over a measurable transformation

defined on a bounded Borel set in a Euclidian space of finite dimension. We next consider (in Section 4) a symbolic representation of the flow $\{P^t\}$ and show that it is, up to an isomorphism, a suspension flow over a countable alphabet topological Markov shift (denote the alphabet by $\mathcal{A}$). Theorem 2.2 cannot yet be applied directly to this suspension flow, since, if for no other reason than that its roof function is not bounded away from zero and has non-summable variations. That is why we change the base (Poincaré section) of our flow for its appropriate cylinder subset (we in fact use a family of cylinders) and thus perform a change-over to a new suspension representation (going back to Veech). The new suspension flow appears to be built over a countable alphabet Bernoulli shift whatever cylinder set we take (this is a Markov–Bernoulli reduction as defined in Subsection 2.3). Not all cylinders are useful for us here, but only those corresponding to admissible “positive” words $w \in \bigcup_{n=1}^{\infty} \mathcal{A}^n$ (there is a canonical way to assign a matrix with non-negative integer entries to each w ; if all the entries are positive, we refer to w as a positive word). If we change the positivity for a stronger requirement that each words w involved in the construction has a simple positive prefix (the definition is given in Subsection 2.1), it will be possible to prove (see Subsections 4.3 and 4.4) that the roof function has summable variations (it is even Hölder continuous), while the measure μ_κ (see Section 1) induces on the base of our suspension flow an invariant measure satisfying the requirements imposed on the measure $\bar{\mu}$ in Theorem 2.2. It remains to note that for each $\{P^t\}$ -invariant ergodic probability measure ν with positive entropy, there is a sufficiently large collection of words w with $\nu(w) > 0$ that have a simple positive prefix (see Lemma 5.5). We thus have everything necessary for applying Theorem 2.2.

3.1 Induction maps

Let π be a permutation of m symbols, which will always be assumed irreducible in the sense that $\pi\{1, \dots, k\} = \{1, \dots, k\}$ implies $k = m$. The Rauzy operations a and b are defined by the formulas

$$a\pi(j) = \begin{cases} \pi j, & \text{if } j \leq \pi^{-1}m, \\ \pi m, & \text{if } j = \pi^{-1}m + 1, \\ \pi(j - 1), & \text{if } \pi^{-1}m + 1 < j \leq m; \end{cases}$$

$$b\pi(j) = \begin{cases} \pi j, & \text{if } \pi j \leq \pi m, \\ \pi j + 1, & \text{if } \pi m < \pi j < m, \\ \pi m + 1, & \text{if } \pi j = m. \end{cases}$$

These operations preserve irreducibility. The *Rauzy class* $\mathcal{R}(\pi)$ is defined as the set of all permutations that can be obtained from π by application of the transformation group generated by a and b . From now on we fix a Rauzy class $\mathcal{R}$ and assume that it consists of irreducible permutations.

For $i, j = 1, \dots, m$, denote by E^{ij} the $m \times m$ matrix whose (i, j) th entry is 1, while all others are zeros. Let E be the identity $m \times m$ -matrix. Following Veech [21], introduce the unimodular matrices

$$A(a, \pi) = \sum_{i=1}^{\pi^{-1}m} E^{ii} + E^{m, \pi^{-1}m+1} + \sum_{i=\pi^{-1}m}^{m-1} E^{i, i+1}, \quad (3.1)$$

$$A(b, \pi) = E + E^{m, \pi^{-1}m}. \quad (3.2)$$

For a vector $\lambda = (\lambda_1, \dots, \lambda_m) \in \mathbb{R}^m$, we write

$$|\lambda| = \sum_{i=1}^m \lambda_i.$$

Let

$$\Delta_{m-1} = \{\lambda \in \mathbb{R}^m : |\lambda| = 1, \lambda_i > 0 \text{ for } i = 1, \dots, m\}.$$

One can identify each pair (λ, π) , $\lambda \in \Delta_{m-1}$, with the *interval exchange map* of the interval $I := [0, 1)$ as follows. Divide I into the sub-intervals $I_k := [\beta_{k-1}, \beta_k)$, where $\beta_0 = 0$, $\beta_k = \sum_{i=1}^k \lambda_i$, $1 \leq k \leq m$, and then place the intervals I_k in I in the following order (from left to right): $I_{\pi^{-1}1}, \dots, I_{\pi^{-1}m}$. We obtain a piecewise linear transformation of I that preserves the Lebesgue measure.

The space $\Delta(\mathcal{R})$ of interval exchange maps corresponding to $\mathcal{R}$ is defined by

$$\Delta(\mathcal{R}) = \Delta_{m-1} \times \mathcal{R}.$$

Denote

$$\Delta_{\pi}^{+} = \{\lambda \in \Delta_{m-1} \mid \lambda_{\pi^{-1}m} > \lambda_m\}, \quad \Delta_{\pi}^{-} = \{\lambda \in \Delta_{m-1} \mid \lambda_m > \lambda_{\pi^{-1}m}\},$$

$$\Delta^{+}(\mathcal{R}) = \cup_{\pi \in \mathcal{R}} \{(\pi, \lambda) \mid \lambda \in \Delta_{\pi}^{+}\},$$

$$\Delta^-(\mathcal{R}) = \cup_{\pi \in \mathcal{R}} \{(\pi, \lambda) \mid \lambda \in \Delta_\pi^-\},$$

$$\Delta^\pm(\mathcal{R}) = \Delta^+(\mathcal{R}) \cup \Delta^-(\mathcal{R}).$$

The *Rauzy-Veech induction map* $\mathcal{T} : \Delta^\pm(\mathcal{R}) \rightarrow \Delta(\mathcal{R})$ is defined as follows:

$$\mathcal{T}(\lambda, \pi) = \begin{cases} (\frac{A(a, \pi)^{-1}\lambda}{|A(a, \pi)^{-1}\lambda|}, a\pi), & \text{if } \lambda \in \Delta_\pi^+, \\ (\frac{A(b, \pi)^{-1}\lambda}{|A(b, \pi)^{-1}\lambda|}, b\pi), & \text{if } \lambda \in \Delta_\pi^-. \end{cases} \quad (3.3)$$

One can check that $\mathcal{T}(\lambda, \pi)$ is the interval exchange map induced by (λ, π) on the interval $[0, 1 - \gamma]$, where $\gamma = \min(\lambda_m, \lambda_{\pi^{-1}m})$; this interval stretches to unit length.

Denote

$$\Delta^\infty(\mathcal{R}) = \bigcap_{n \geq 0} \mathcal{T}^{-n} \Delta^\pm(\mathcal{R}). \quad (3.4)$$

Every $\mathcal{T}$ -invariant probability measure is concentrated on $\Delta^\infty(\mathcal{R})$. On the other hand, a natural Lebesgue measure defined on $\Delta(\mathcal{R})$, which is finite, but non-invariant, is also concentrated on $\Delta^\infty(\mathcal{R})$. Veech [21] showed that $\mathcal{T}$ has an absolutely continuous ergodic invariant measure on $\Delta(\mathcal{R})$, which is, however, infinite.

Following Zorich [26], for $(\lambda, \pi) \in \Delta^\infty(\mathcal{R})$, we set

$$n(\lambda, \pi) = \begin{cases} \min\{k > 0 : \mathcal{T}^k(\lambda, \pi) \in \Delta^-(\mathcal{R})\}, & \text{if } \lambda \in \Delta_\pi^+; \\ \min\{k > 0 : \mathcal{T}^k(\lambda, \pi) \in \Delta^+(\mathcal{R})\}, & \text{if } \lambda \in \Delta_\pi^-. \end{cases} \quad (3.5)$$

The *Rauzy-Veech-Zorich induction map* $\mathcal{G}$ is defined by the formula

$$\mathcal{G}(\lambda, \pi) = \mathcal{T}^{n(\lambda, \pi)}(\lambda, \pi), \quad (\lambda, \pi) \in \Delta^\infty(\mathcal{R}). \quad (3.6)$$

Theorem 3.1 (Zorich [26]). *The map $\mathcal{G}$ has an ergodic invariant probability measure ν absolutely continuous with respect to the Lebesgue measure on $\Delta^\infty(\mathcal{R})$. The density ρ of this measure is of the form*

$$\rho(\lambda, \pi) = P_\pi(\lambda)/Q_\pi(\lambda), \quad \lambda = (\lambda_1, \dots, \lambda_m),$$

where P_π and Q_π are homogeneous polynomials with non-negative coefficients.

3.2 Zippered rectangles

Here we briefly recall the construction of the Veech space of zippered rectangles. We use the notation of [4].

Zippered rectangles associated with the Rauzy class $\mathcal{R}$ are triples (λ, π, δ) , where $\lambda = (\lambda_1, \dots, \lambda_m) \in \mathbb{R}^m$, $\lambda_i > 0$, $\pi \in \mathcal{R}$, $\delta = (\delta_1, \dots, \delta_m) \in \mathbb{R}^m$, and the vector δ satisfies the following inequalities:

$$\delta_1 + \dots + \delta_i \leq 0, \quad i = 1, \dots, m-1, \quad (3.7)$$

$$\delta_{\pi^{-1}1} + \dots + \delta_{\pi^{-1}i} \geq 0, \quad i = 1, \dots, m-1. \quad (3.8)$$

The set of all vectors δ satisfying (3.7), (3.8) is a cone in $\mathbb{R}^m$; we denote it by $K(\pi)$.

For a zippered rectangle (λ, π, δ) , we set

$$h_r := -\sum_{i=1}^{r-1} \delta_i + \sum_{i=1}^{\pi r-1} \delta_{\pi^{-1}i}, \quad (3.9)$$

$$Area(\lambda, \pi, \delta) := \sum_{r=1}^m \lambda_r h_r. \quad (3.10)$$

(Our convention is $\sum_{i=u}^v \dots = 0$ when $u > v$.) By (3.7), (3.8) $h_r \geq 0$ for all r , and if we relate the set $\mathcal{Z} := \bigcup_{r=1}^m I_r \times [0, h_r]$ (a union of rectangles in $\mathbb{R}^2$) to every triple (λ, π, δ) , then $Area(\lambda, \pi, \delta)$ becomes merely the Lebesgue measure (area) of $\mathcal{Z}$. By appropriate identification of intervals in the boundaries of different rectangles $I_r \times [0, h_r]$ one obtains a compact Riemannian surface and a 1-form on it. This procedure (due to Veech [21]) is thoroughly described in the literature (see, for example, [25], [26]). We do not use it as such and so omit details.

Denote by $\mathcal{V}(\mathcal{R})$ the space of all zippered rectangles corresponding to the Rauzy class $\mathcal{R}$, i.e.,

$$\mathcal{V}(\mathcal{R}) = \{(\lambda, \pi, \delta) : \lambda \in \mathbb{R}_+^m, \pi \in \mathcal{R}, \delta \in K(\pi)\}.$$

Let also

$$\mathcal{V}^+(\mathcal{R}) = \{(\lambda, \pi, \delta) \in \mathcal{V}(\mathcal{R}) : \lambda_{\pi^{-1}m} > \lambda_m\},$$

$$\mathcal{V}^-(\mathcal{R}) = \{(\lambda, \pi, \delta) \in \mathcal{V}(\mathcal{R}) : \lambda_{\pi^{-1}m} < \lambda_m\},$$

$$\mathcal{V}^\pm(\mathcal{R}) = \mathcal{V}^+(\mathcal{R}) \cup \mathcal{V}^-(\mathcal{R}).$$

Veech [21] introduced the flow $\{P^t\}$ acting on $\mathcal{V}(\mathcal{R})$ by the formula

$$P^t(\lambda, \pi, \delta) = (e^t \lambda, \pi, e^{-t} \delta),$$

and the map $\mathcal{U} : \mathcal{V}^\pm(\mathcal{R}) \rightarrow \mathcal{V}(\mathcal{R})$, where

$$\mathcal{U}(\lambda, \pi, \delta) = \begin{cases} (A(\pi, a)^{-1} \lambda, a\pi, A(\pi, a)^{-1} \delta), & \text{if } \lambda_{\pi^{-1}m} > \lambda_m, \\ (A(\pi, b)^{-1} \lambda, b\pi, A(\pi, b)^{-1} \delta), & \text{if } \lambda_{\pi^{-1}m} < \lambda_m. \end{cases}$$

(The inclusion $\mathcal{U}\mathcal{V}^\pm(\mathcal{R}) \subset \mathcal{V}(\mathcal{R})$ is not evident and should be proved; this was done in [21].) The map $\mathcal{U}$ and the flow $\{P^t\}$ commute on $\mathcal{V}^\pm(\mathcal{R})$ and both preserve the measure determined on $\mathcal{V}(\mathcal{R})$ by the volume form $Vol = d\lambda_1 \dots d\lambda_m d\delta_1 \dots d\delta_m$. They also preserve the area of a zippered rectangle (see (3.10)) and hence can be restricted to the set

$$\mathcal{V}^{1,\pm}(\mathcal{R}) := \{(\lambda, \pi, \delta) \in \mathcal{V}^\pm(\mathcal{R}) : Area(\lambda, \pi, \delta) = 1\}.$$

The restriction of the volume form Vol to $\mathcal{V}^{1,\pm}(\mathcal{R})$ induces on this set a measure $\mu_{\mathcal{R}}$ which is invariant under $\mathcal{U}$ and $\{P^t\}$.

For $(\lambda, \pi) \in \Delta(\mathcal{R})$, denote

$$\tau^0(\lambda, \pi) =: -\log(|\lambda| - \min(\lambda_m, \lambda_{\pi^{-1}m})). \quad (3.11)$$

From (3.1), (3.2) it follows that if $\lambda \in \Delta_\pi^+ \cup \Delta_\pi^-$, then

$$\tau^0(\lambda, \pi) = -\log |A^{-1}(c, \pi)\lambda|, \quad (3.12)$$

where $c = a$ when $\lambda \in \Delta_\pi^+$, and $c = b$ when $\lambda \in \Delta_\pi^-$.

Next denote

$$\mathcal{Y}_1(\mathcal{R}) := \{x = (\lambda, \pi, \delta) \in \mathcal{V}(\mathcal{R}) : |\lambda| = 1, Area(\lambda, \pi, \delta) = 1\},$$

$$\tau(x) := \tau^0(\lambda, \pi) \text{ for } x = (\lambda, \pi, \delta) \in \mathcal{Y}_1(\mathcal{R}),$$

$$\mathcal{V}_{1,\tau}(\mathcal{R}) := \bigcup_{x \in \mathcal{Y}_1(\mathcal{R}), 0 \leq t \leq \tau(x)} P^t x. \quad (3.13)$$

Using the map $\mathcal{U}$, we are going to transfer the flow P^t to the set $\mathcal{V}_{1,\tau}(\mathcal{R})$ (or, more precisely, to its proper subset).

It is easy to check that $\mathcal{U}P^{\tau(x)}x \in \mathcal{Y}_1(\mathcal{R})$ for every $x \in \mathcal{Y}_1(\mathcal{R}) \cap \mathcal{V}^\pm(\mathcal{R})$. Identifying the points $P^{\tau(x)}x$ and $\mathcal{U}P^{\tau(x)}x$, we can continue the trajectory of

x by some distance. But it can happen that $\mathcal{U}P^{\tau(x)}x \notin \mathcal{V}^{\pm}(\mathcal{R})$, so that we cannot proceed this way.

To make $\{P^t\}$ well defined on an invariant set we have to somewhat reduce the domain of $\mathcal{U}$. Let

$$\mathcal{V}_{\neq}^{1,\pm}(\mathcal{R}) := \{(\lambda, \pi, \delta) \in \mathcal{V}^{1,\pm}(\mathcal{R}) : a_m(\delta) \neq 0\},$$

$$\mathcal{V}_{\infty}(\mathcal{R}) := \bigcap_{n \in \mathbb{Z}} \mathcal{U}^n \mathcal{V}_{\neq}^{1,\pm}(\mathcal{R}).$$

Clearly $\mathcal{U}^n$ is well-defined on $\mathcal{V}_{\infty}(\mathcal{R})$ for all $n \in \mathbb{Z}$.

We now set

$$\mathcal{Y}(\mathcal{R}) := \mathcal{Y}_1(\mathcal{R}) \cap \mathcal{V}_{\infty}(\mathcal{R}), \quad \tilde{\mathcal{V}}(\mathcal{R}) := \mathcal{V}_{1,\tau}(\mathcal{R}) \cap \mathcal{V}_{\infty}(\mathcal{R}).$$

The above identification enables us to define on $\tilde{\mathcal{V}}(\mathcal{R})$ a natural flow, for which we retain the notation $\{P^t\}$. (Although the bounded positive function τ is not separated from zero, the flow $\{P^t\}$ is well defined.)

Let us also note that $\mu_{\mathcal{R}}(\mathcal{V}_{1,\tau}(\mathcal{R})) > 0$. By a theorem proven independently by Veech [21] and Masur [15] $\mu_{\mathcal{R}}(\mathcal{V}_{1,\tau}(\mathcal{R})) < \infty$, and we in what follows assume that the restriction of $\mu_{\mathcal{R}}$ to $\mathcal{V}_{1,\tau}(\mathcal{R})$ is normalized to have total mass 1. Since clearly $\mu_{\mathcal{R}}(\mathcal{V}_{1,\tau}(\mathcal{R}) \setminus \tilde{\mathcal{V}}(\mathcal{R})) = 0$, we can consider $\mu_{\mathcal{R}}$ to be defined on $\tilde{\mathcal{V}}(\mathcal{R})$. This measure is ergodic with respect to the flow $\{P^t\}$.

Remark 3.2. The presentation here differs from the one in Veech [21] by a linear change of variable: the Veech vector parameters h and a of a zippered rectangle (λ, π, δ) are expressed in terms of π and δ by (3.9) and the equations

$$a_r = a_r(\delta) = - \sum_{i=1}^r \delta_i, \quad r = 1, \dots, m.$$

Following Zorich [26], denote

$$\mathcal{Y}^+(\mathcal{R}) = \{x = (\lambda, \pi, \delta) \in \mathcal{Y}(\mathcal{R}) : \lambda \in \Delta_{\pi}^+, a_m(\delta) < 0\},$$

$$\mathcal{Y}^-(\mathcal{R}) = \{x = (\lambda, \pi, \delta) \in \mathcal{Y}(\mathcal{R}) : \lambda \in \Delta_{\pi}^-, a_m(\delta) > 0\},$$

$$\mathcal{Y}^{\pm}(\mathcal{R}) = \mathcal{Y}^+(\mathcal{R}) \cup \mathcal{Y}^-(\mathcal{R}),$$

and let $\mathcal{V}_{\infty}^{\pm}(\mathcal{R})$ be the set of all $x \in \mathcal{Y}^{\pm}(\mathcal{R})$ for which there exist infinitely many positive t and infinitely many negative t such that $P^t x \in \mathcal{Y}^{\pm}(\mathcal{R})$.

Take $x \in \mathcal{Y}_\infty^\pm(\mathcal{R})$, $x = (\lambda, \pi, \delta)$, and let $\mathcal{F}(x)$ be the first return of x to the transversal $\mathcal{Y}^\pm(\mathcal{R})$ under the flow $\{P^t\}$. The map $\mathcal{F}$ is an extension of the map $\mathcal{G}$ to the space of zippered rectangles:

$$\text{if } \mathcal{F}(\lambda, \pi, \delta) = (\lambda', \pi', \delta'), \text{ then } (\lambda', \pi') = \mathcal{G}(\lambda', \pi'). \quad (3.14)$$

Note that $\mathcal{F}$ is invertible on $\mathcal{Y}_\infty^\pm(\mathcal{R})$ and if $x \in \mathcal{Y}^+(\mathcal{R})$ (respectively, $x \in \mathcal{Y}^-(\mathcal{R})$), then $\mathcal{F}(x) \in \mathcal{Y}^-(\mathcal{R})$ (respectively, $\mathcal{F}(x) \in \mathcal{Y}^+(\mathcal{R})$). Moreover,

$$\mathcal{Y}_\infty^\pm(\mathcal{R}) = \bigcap_{n \in \mathbb{Z}} \mathcal{F}^n \mathcal{Y}^\pm(\mathcal{R}).$$

If $x = (\lambda, \pi, \delta) \in \mathcal{Y}_\infty^\pm(\mathcal{R})$ and $\mathcal{G}(\lambda, \pi) = \mathcal{T}^n(\lambda, \pi)$, then, by (3.3), (3.6), the first return time of x to $\mathcal{Y}^\pm(\mathcal{R})$ under the flow $\{P^t\}$ is

$$\tau(\lambda, \pi) + \dots + \tau(\mathcal{T}^{n-1}(\lambda, \pi)) = -\log |A^{-1}(c, c^{n-1}\pi) \dots A^{-1}(c, \pi)\lambda|, \quad (3.15)$$

where $c = a$ when $\lambda \in \Delta_\pi^+$, and $c = b$ when $\lambda \in \Delta_\pi^-$.

We finish this section with considering a relationship between the probability measure $\mu_{\mathcal{R}}$ mentioned above and the measure ν introduced in Theorem 3.1.

Denote by $\mu_{\mathcal{R}}^1$ the $\mathcal{F}$ -invariant probability measure induced by $\mu_{\mathcal{R}}$ on $\mathcal{Y}_\infty^\pm(\mathcal{R})$. Remark that if $(\lambda, \pi, \delta) \in \mathcal{Y}_\infty^\pm(\mathcal{R})$, then $(\lambda, \pi) \in \Delta^\infty$.

Lemma 3.3. (cf. [21], [26]) *Let $\tilde{\psi} : \mathcal{Y}_\infty^\pm(\mathcal{R}) \rightarrow \Delta^\infty(\mathcal{R})$ be the map defined by $\tilde{\psi}(\lambda, \pi, \delta) = (\lambda, \pi)$. Then $\tilde{\psi}_* \mu_{\mathcal{R}}^1 = \nu$.*

Proof. Note that there is a natural Lebesgue measure on each of the spaces $\tilde{\mathcal{V}}(\mathcal{R})$, $\mathcal{Y}_\infty^\pm(\mathcal{R})$, and $\Delta^\infty(\mathcal{R})$. Since $\mu_{\mathcal{R}}$ is proportional to the Lebesgue measure on $\tilde{\mathcal{V}}(\mathcal{R})$, from the definition of $\{P^t\}$ it follows that $\mu_{\mathcal{R}}^1$ is absolutely continuous with respect to the Lebesgue measure on $\mathcal{Y}_\infty^\pm(\mathcal{R})$. Let $\nu^1 := \tilde{\psi}_* \mu_{\mathcal{R}}^1$. It is clear that ν^1 is a probability measure absolutely continuous with respect to mes_Δ , the Lebesgue measure on $\Delta^\infty(\mathcal{R})$, while by Theorem 3.1 the probability measure ν is equivalent to mes_Δ and ergodic with respect to $\mathcal{G}$. Therefore, $\nu^1 = \nu$. $\square$

4 Symbolic representation of the covering flow

In this section we construct suspension flows over symbolic Markov shifts that will be of great importance in the study of the flow $\{P^t\}$. Using [21], [26], we begin with a brief description of a symbolic model for the map $\mathcal{G}$.

4.1 Symbolic dynamics for the mapping $\mathcal{G}$

We shall only deal with the interval exchanges (λ, π) from $\Delta^\infty(\mathcal{R})$ (see (3.4)), so that all iterations of the map $\mathcal{G}$ are defined. Our notation follows [4].

Consider the alphabet

$$\mathcal{A} := \{(c, n, \pi) \mid c = a \text{ or } b, n \in \mathbb{N}, \pi \in \mathcal{R}\}.$$

For $w_1 = (c_1, n_1, \pi_1) \in \mathcal{A}$, $w_2 = (c_2, n_2, \pi_2) \in \mathcal{A}$, we set

$$B(w_1, w_2) = \begin{cases} 1 & \text{if } (c_1)^{n_1} \pi_1 = \pi_2, c_2 \neq c_1, \\ 0 & \text{otherwise} \end{cases}$$

and thus define a function $B : \mathcal{A} \times \mathcal{A} \rightarrow \{0, 1\}$. In other terms, we have a directed graph $\Gamma_{\mathcal{A}, B} = (V, E)$, where $V = \mathcal{A}$ and where $(w_1, w_2) \in E$ if and only if $B(w_1, w_2) = 1$. From the definition of the Rauzy class $\mathcal{R}$ in Subsection 3.1 it follows that the graph $\Gamma_{\mathcal{A}, B}$ is connected.

Introduce the space of words

$$\mathcal{W}_{\mathcal{A}, B} = \{w = w_1 \dots w_n \mid w_i \in \mathcal{A}, B(w_i, w_{i+1}) = 1 \text{ for } i = 1, \dots, n\}.$$

It is convenient to include the empty word in $\mathcal{W}_{\mathcal{A}, B}$. We use the notation of Sub-section 2.1. In particular, for a word $w \in \mathcal{W}_{\mathcal{A}, B}$, we denote by $|w|$ its length, i.e., the number of symbols in it; given two words $w^{(1)}, w^{(2)} \in \mathcal{W}_{\mathcal{A}, B}$, we denote by $w^{(1)}w^{(2)}$ their concatenation. Note that the word $w^{(1)}w^{(2)}$ need not belong to $\mathcal{W}_{\mathcal{A}, B}$, unless a compatibility condition is satisfied by the last letter of $w^{(1)}$ and the first letter of $w^{(2)}$.

To each nonempty word $w \in \mathcal{W}_{\mathcal{A}, B}$ we assign a *renormalization matrix* $A(w)$ as follows. If w is a single-letter word, $w = (c, n, \pi) \in \mathcal{A}$, we set (see (3.1), (3.2))

$$A(w) = A(c, \pi)A(c, c\pi) \dots A(c, c^{n-1}\pi); \quad (4.1)$$

for $w \in \mathcal{W}_{\mathcal{A}, B}$, where $w = w_1 \dots w_n$, $w_i \in \mathcal{A}$, we set

$$A(w) = A(w_1) \dots A(w_n). \quad (4.2)$$

Consider the sequence spaces

$$\Omega_{\mathcal{A}, B} = \{\omega = (\omega_0, \omega_1, \dots) \mid \omega_n \in \mathcal{A}, B(\omega_n, \omega_{n+1}) = 1 \text{ for all } n \in \mathbb{Z}_+\},$$

and

$$\Omega_{\mathcal{A},B}^{\mathbb{Z}} = \{\omega = (\dots, \omega_{-1}, \omega_0, \omega_1, \dots) \mid \omega_n \in \mathcal{A}, B(\omega_n, \omega_{n+1}) = 1 \text{ for all } n \in \mathbb{Z}\}.$$

Denote by σ the one step left shift on both these spaces.

Let us now describe the coding map. For every letter $w = (c, n, \pi) \in \mathcal{A}$, we set

$$\Delta(w) = \begin{cases} \Delta^+(\mathcal{R}) \cap \{(\lambda, \pi) \in \Delta^\infty(\mathcal{R}) \mid n(\lambda, \pi)\} & \text{if } c = a, \\ \Delta^-(\mathcal{R}) \cap \{(\lambda, \pi) \in \Delta^\infty(\mathcal{R}) \mid n(\lambda, \pi)\} & \text{if } c = b. \end{cases} \quad (4.3)$$

In other words, when $c = a$ (resp., $c = b$), $\Delta(w)$ consists of all points $(\lambda, \pi) \in \Delta^\infty(\mathcal{R}) \cap \Delta^+(\mathcal{R})$ (resp., $(\lambda, \pi) \in \Delta^\infty(\mathcal{R}) \cap \Delta^-(\mathcal{R})$) such that $\mathcal{T}^k(\lambda, \pi) \in \Delta^+(\mathcal{R})$ (resp., $\mathcal{T}^k(\lambda, \pi) \in \Delta^-(\mathcal{R})$) for $k = 0, \dots, n-1$, and $\mathcal{T}^n(\lambda, \pi) \in \Delta^-(\mathcal{R})$ (resp., $\mathcal{T}^k(\lambda, \pi) \in \Delta^+(\mathcal{R})$). Using (3.5), one also can check that if $w = (c, n, \pi)$, then

$$\Delta(w) = \begin{cases} \Delta^\infty(\mathcal{R}) \cap \{(\lambda, \pi) \mid \lambda \in \Delta_\pi^+, \frac{A(w)^{-1}\lambda}{|A(w)^{-1}\lambda|} \in \Delta_{a^n\pi}^-\}, & \text{if } c = a, \\ \Delta^\infty(\mathcal{R}) \cap \{(\lambda, \pi) \mid \lambda \in \Delta_\pi^-, \frac{A(w)^{-1}\lambda}{|A(w)^{-1}\lambda|} \in \Delta_{b^n\pi}^+\}, & \text{if } c = b. \end{cases} \quad (4.4)$$

It is easy to see that all the sets $\Delta(w)$, $w \in \mathcal{A}$, are non-empty and constitute a partition of $\Delta^\infty(\mathcal{R})$. We use this partition to construct a symbolic dynamics for $\mathcal{G}$. By iterating this partition n times under the action of the transformation $\mathcal{G}$ we obtain a partition whose elements, $\Delta(w)$, are determined by the words $w \in \mathcal{W}_{\mathcal{A},B}$ of length n . Namely, for a word $w = w_1 \dots w_n \in \mathcal{W}_{\mathcal{A},B}$, $w_i \in \mathcal{A}$, we set

$$\Delta(w) = \bigcap_{i=0}^{n-1} \mathcal{G}^{-i} \Delta(w_{i+1}). \quad (4.5)$$

Remark 4.1. From (3.3), (3.6), and (4.1)–(4.4) it follows that if $(\lambda, \pi) \in \Delta(w)$ and $(\lambda', \pi') = \mathcal{G}(\lambda, \pi)$, then

$$\lambda' = A^{-1}(w)\lambda / |A^{-1}(w)\lambda|, \quad \lambda = A(w)\lambda' / |A(w)\lambda'|.$$

These formulas can be easily extended by induction to the case where $w = w_1 \dots w_n \in \mathcal{W}_{\mathcal{A},B}$, $(\lambda, \pi) \in \Delta(w)$, and $(\lambda', \pi') = \mathcal{G}(\lambda, \pi)$.

The *coding map* $\Phi : \Delta^\infty(\mathcal{R}) \rightarrow \Omega_{\mathcal{A},B}$ is given by the formula

$$\Phi(\lambda, \pi) = (\omega_0, \omega_1, \dots) \text{ if } \mathcal{G}^n(\lambda, \pi) \in \Delta(\omega_n) \text{ for } n = 0, 1, \dots \quad (4.6)$$

Consider conditions under which the coding map is invertible. For $\mathbf{q} \in \mathcal{W}_{\mathcal{A},B}$ we denote by $\Omega_{\mathbf{q}}$ the set of all sequences $\omega \in \Omega_{\mathcal{A},B}$ starting from the word $\mathbf{q}$ and containing infinitely many occurrences of $\mathbf{q}$. A key role will be played below by the words $\mathbf{q}$ such that all entries of the renormalization matrix $A(\mathbf{q})$ are positive. For short we will refer to these $\mathbf{q}$ as *positive words*. Observe that each word containing a positive prefix is also positive.

The next two lemmas are due to Veech [21].

Lemma 4.2. *Let $\mathbf{q} \in \mathcal{W}_{\mathcal{A},B}$ be a positive word. Then for every $\omega \in \Omega_{\mathbf{q}}$, there exists a unique $(\lambda, \pi) \in \Delta^\infty(\mathcal{R})$ such that $\Phi(\lambda, \pi) = \omega$.*

Lemma 4.3. *If the interval exchange map $(\lambda, \pi) : [0, 1] \rightarrow [0, 1]$ is uniquely ergodic, then there exists a positive word $\mathbf{q} \in \mathcal{W}_{\mathcal{A},B}$ such that $(\lambda, \pi) \in \Delta(\mathbf{q})$.*

Remark 4.4. Veech [21] in fact observed that for the existence of $\mathbf{q}$ mentioned in Lemma 4.3 it is sufficient for (λ, π) to satisfy Keane's infinite distinct orbit condition.

4.2 Symbolic dynamics for the flow $\{P^t\}$

We first construct a symbolic dynamics for the map $\mathcal{F}$ introduced in Subsection 3.2. For $(\lambda, \pi, \delta) \in \mathcal{Y}_\infty^\pm(\mathcal{R})$ we set

$$\tilde{\Phi}(\lambda, \pi, \delta) = (\dots, \omega_{-1}, \omega_0, \omega_1, \dots), \quad \omega_i \in \mathcal{A}, \quad (4.7)$$

if $\mathcal{F}^n(\lambda, \pi, \delta) = (\lambda'_n, \pi'_n, \delta'_n)$ and $(\lambda'_n, \pi'_n) \in \Delta(\omega_n)$, $n \in \mathbb{Z}$ (remind that $\mathcal{F}$ is invertible on $\mathcal{Y}_\infty^\pm(\mathcal{R})$). In parallel with the coding map Φ (see Subsection 4.1) we have

$$\tilde{\Phi}(\mathcal{Y}_\infty^\pm(\mathcal{R})) \subset \Omega_{\mathcal{A},B}^\mathbb{Z}.$$

Moreover, from (3.14) and (4.6) it follows that (4.7) implies that

$$\Phi(\lambda, \pi) = (\omega_0, \omega_1, \dots).$$

For $\mathbf{q} \in \mathcal{W}_{\mathcal{A},B}$, $|\mathbf{q}| = l$, similarly to the definition of $\Omega_{\mathbf{q}}$ in Subsection 4.1, we denote by $\Omega_{\mathbf{q}}^\mathbb{Z}$ the set of all sequences $\omega \in \Omega_{\mathcal{A},B}^\mathbb{Z}$ satisfying $\omega_0 \dots \omega_{l-1} = \mathbf{q}$ and admitting infinitely many occurrences, both in the past and in the future, of the word $\mathbf{q}$.

Let

$$\mathcal{Y}_{\mathbf{q},\infty}^\pm(\mathcal{R}) := \tilde{\Phi}^{-1}(\Omega_{\mathbf{q}}^\mathbb{Z}), \quad \mathcal{V}_{\mathbf{q}}(\mathcal{R}) := \bigcup_{t \in \mathbb{R}} P^t \mathcal{Y}_{\mathbf{q},\infty}^\pm(\mathcal{R}),$$

and assume that $\mathcal{Y}_{\mathbf{q},\infty}^{\pm}(\mathcal{R})$ (and hence $\mathcal{V}_{\mathbf{q}}(\mathcal{R})$) is non-empty.

Let $\mathcal{F}_{\mathbf{q}}$ be the first return map of $\mathcal{F}$ to $\mathcal{Y}_{\mathbf{q},\infty}^{\pm}(\mathcal{R})$, i.e., the map induced by $\mathcal{F}$ on $\mathcal{Y}_{\mathbf{q},\infty}^{\pm}(\mathcal{R})$ (cf. Subsection 2.3).

By definition, $\{P^t\}|_{\mathcal{V}_{\mathbf{q}}(\mathcal{R})}$, the restriction of the flow $\{P^t\}$ to $\mathcal{V}_{\mathbf{q}}(\mathcal{R})$, is Borel isomorphic to a suspension flow $(\mathcal{F}_{\mathbf{q}}, \tau_{\mathbf{q}})$ over the map $\mathcal{F}_{\mathbf{q}}$. To describe the roof function $\tau_{\mathbf{q}}$ we take $(\lambda, \pi) \in \Delta^{\infty}(\mathcal{R})$, $(\lambda', \pi') = \mathcal{G}(\lambda, \pi)$, and introduce, following Veech, the function

$$\tau^1 : (\lambda, \pi) \mapsto \log |A(\omega_0)\lambda'|, \quad (4.8)$$

where ω_0 is determined by the equation $\Phi(\lambda, \pi) = (\omega_0, \omega_1, \dots)$, that is $(\lambda, \pi) \in \Delta(\omega_0)$. (Recall that the norm of a vector v is given by $|v| = \sum_i |v_i|$.) Using (3.14), (3.15), it is easy to check that if $x = (\lambda, \pi, \delta) \in \mathcal{Y}_{\infty}^{\pm}(\mathcal{R})$ and $(\lambda, \pi) \in \Delta(\omega_0)$, where $\omega_0 = (c, n, \pi)$, then the first return time of x to $\mathcal{Y}_{\infty}^{\pm}(\mathcal{R})$ under the action of the flow $\{P^t\}$ is just $\tau^1(\lambda, \pi)$.

Let now $(\lambda, \pi) \in \Phi^{-1}(\Omega_{\mathbf{q}})$, $(\omega_0, \omega_1, \dots) = \Phi(\lambda, \pi)$, and let s be the moment of the second appearance of the word $\mathbf{q}$ in $(\omega_0, \omega_1, \dots)$, i.e.,

$$s = s(\omega_0, \omega_1, \dots) = \min\{k > 0 \mid (\omega_k, \dots, \omega_{k+l-1}) = \mathbf{q}\}. \quad (4.9)$$

Denote

$$\tau_{\mathbf{q}}^1(\lambda, \pi) = \tau^1(\lambda, \pi) + \tau^1(\mathcal{G}(\lambda, \pi)) + \dots + \tau^1(\mathcal{G}^{s-1}(\lambda, \pi)). \quad (4.10)$$

If $x = (\lambda, \pi, \delta) \in \mathcal{Y}_{\mathbf{q},\infty}^{\pm}(\mathcal{R})$, then $(\lambda, \pi) \in \Phi^{-1}(\Omega_{\mathbf{q}})$, and we have

$$\tau_{\mathbf{q}}(x) = \tau_{\mathbf{q}}^1(\lambda, \pi). \quad (4.11)$$

Denote by Ψ_1 the map from $\mathcal{V}_{\mathbf{q}}(\mathcal{R})$ to the phase space of $(\mathcal{F}_{\mathbf{q}}, \tau_{\mathbf{q}})$ that induces the above-mentioned isomorphism between the flows $\{P^t\}|_{\mathcal{V}_{\mathbf{q}}(\mathcal{R})}$ and $(\mathcal{F}_{\mathbf{q}}, \tau_{\mathbf{q}})$.

From now on we assume that $\mathbf{q}$ is a positive word. For such $\mathbf{q}$, we construct a suspension flow $(\hat{\sigma}_{\mathbf{q}}, \hat{\tau}_{\mathbf{q}})$ closely related to $(\mathcal{F}_{\mathbf{q}}, \tau_{\mathbf{q}})$. Denote, as before, by σ the one-step left shift on $\Omega_{\mathcal{A},B}^{\mathbb{Z}}$ and, for $\omega = (\dots, \omega_{-1}, \omega_0, \omega_1, \dots) \in \Omega_{\mathbf{q}}^{\mathbb{Z}}$, let

$$\hat{\sigma}_{\mathbf{q}}(\omega) := \sigma^s, \quad \hat{\tau}_{\mathbf{q}}(\omega) := \tau_{\mathbf{q}}^1(\Phi^{-1}(\omega_0, \omega_1, \dots)), \quad (4.12)$$

where s is defined in (4.9). (Observe that $\omega \in \Omega_{\mathbf{q}}^{\mathbb{Z}}$ implies $(\omega_0, \omega_1, \dots) \in \Omega_{\mathbf{q}}$, hence by Lemma 4.2 $\Phi^{-1}(\omega_0, \omega_1, \dots)$ is a uniquely defined point from $\Delta^{\infty}(\mathcal{R})$).

Proposition 6 in [4] states that if $\mathbf{q} \in \mathcal{W}_{\mathcal{A},B}$ is positive, then for every $\omega \in \Omega_{\mathbf{q}}^{\mathbb{Z}}$, there exists at most one zippered rectangle corresponding to it; in other words, the map $\tilde{\Phi}$ restricted to the set $\mathcal{Y}_{\mathbf{q},\infty}^{\pm}(\mathcal{R}) := \tilde{\Phi}^{-1}(\Omega_{\mathbf{q}}^{\mathbb{Z}})$ is injective. It induces, in a natural way, a measurable injective map Ψ_2 from the phase space of the flow $(\sigma_{\mathbf{q}}, \tau_{\mathbf{q}})$ to the phase space of the flow $(\hat{\sigma}_{\mathbf{q}}, \hat{\tau}_{\mathbf{q}})$ that sends the former flow to the latter one restricted to some invariant set. Hence $(\mathcal{F}_{\mathbf{q}}, \tau_{\mathbf{q}})$ is embedded (in the sense of Section 2) into $(\hat{\sigma}_{\mathbf{q}}, \hat{\tau}_{\mathbf{q}})$.

Introduce a new alphabet $\mathcal{A}_{\mathbf{q}}$; it will consist of all words $w = (v_1 \dots v_n) \in \mathcal{W}_{\mathcal{A},B}$, $v_i \in \mathcal{A}$, $n > l$, such that $(v_1, \dots, v_l) = \mathbf{q}$, $(v_{n-l+1}, \dots, v_n) = \mathbf{q}$, and no other subword of w coincides with $\mathbf{q}$. Since $|\mathcal{A}| = \infty$ and the graph $\Gamma_{\mathcal{A},B}$ is connected (see Subsection 4.1), we have $|\mathcal{A}_{\mathbf{q}}| = \infty$. By the Markov-Bernoulli reduction, introduced in Subsection 2.3, there is a measurable one-to-one map $\Psi_{M-B} : \Omega_{\mathbf{q}}^{\mathbb{Z}} \rightarrow (\mathcal{A}_{\mathbf{q}})^{\mathbb{Z}}$ that sends $\hat{\sigma}_{\mathbf{q}}$ to the one-step left shift $\sigma_{\mathbf{q}}$ on $(\mathcal{A}_{\mathbf{q}})^{\mathbb{Z}}$. Hence the flow $(\hat{\sigma}_{\mathbf{q}}, \hat{\tau}_{\mathbf{q}})$ is isomorphic to the suspension flow $(\sigma_{\mathbf{q}}, f_{\mathbf{q}})$, where

$$f_{\mathbf{q}}(u) := \hat{\tau}_{\mathbf{q}}(\Psi_{M-B}^{-1}(u)), \quad u \in (\mathcal{A}_{\mathbf{q}})^{\mathbb{Z}}. \quad (4.13)$$

Denote by Ψ_3 the corresponding map from the phase space of $(\hat{\sigma}_{\mathbf{q}}, \hat{\tau}_{\mathbf{q}})$ to the phase space of $(\sigma_{\mathbf{q}}, f_{\mathbf{q}})$. Summing up, we can state the following.

Lemma 4.5. *The mapping $\Psi := \Psi_1 \circ \Psi_2 \circ \Psi_3$ yields an embedding of the flow $\{P^t\}|_{\mathcal{V}_{\mathbf{q}}(\mathcal{R})}$ into the flow $(\sigma_{\mathbf{q}}, f_{\mathbf{q}})$.*

Let us turn to the probability measure $\mu_{\mathcal{R}}$ on $\tilde{\mathcal{V}}(\mathcal{R})$ and the probability measure $\mu_{\mathcal{R}}^1$ induced by $\mu_{\mathcal{R}}$ on $\mathcal{Y}_{\infty}^{\pm}(\mathcal{R})$ (see Subsection 3.2).

Observe that $\mu_{\mathcal{R}}$ assigns a positive mass to every Borel set with nonempty interior. (We assume a natural topology on $\tilde{\mathcal{V}}(\mathcal{R})$ as well as on other spaces we encounter in this paper.) From this fact, using Lemma 4.2 and the definition of $\mathcal{T}$, $\mathcal{G}$, and $\mathcal{F}$, we easily derive that

$$\mu_{\mathcal{R}}^1(\{x \in \mathcal{Y}_{\infty}^{\pm}(\mathcal{R}) : x = (\lambda, \pi, \delta), (\lambda, \pi) \in \Delta(\mathbf{q})\}) > 0.$$

Since $\mu_{\mathcal{R}}$ is $\{P^t\}$ -ergodic, we have

$$\mu_{\mathcal{R}}(\mathcal{V}_{\mathbf{q}}(\mathcal{R})) = 1, \quad \mu_{\mathcal{R}}^1\left(\bigcup_{n \in \mathbb{Z}} \mathcal{F}^n \mathcal{Y}_{\mathbf{q},\infty}^{\pm}(\mathcal{R})\right) = 1. \quad (4.14)$$

By normalizing the restriction of $\mu_{\mathcal{R}}^1$ to $\mathcal{Y}_{\mathbf{q},\infty}^{\pm}(\mathcal{R})$ we obtain a probability measure $\bar{\mu}_{\mathcal{R},\mathbf{q}}^1$.

Let ψ be the natural projection of $\Omega_{\mathcal{A},B}^{\mathbb{Z}}$ on $\Omega_{\mathcal{A},B}$. Recall that in Subsection 3.2 we introduced the natural projection $\tilde{\psi}$ of $\mathcal{Y}_{\infty}^{\pm}(\mathcal{R})$ on $\Delta^{\infty}(\mathcal{R})$. It is clear that $\psi(\tilde{\Phi}(x)) = \Phi(\tilde{\psi}(x))$ for every $x \in \bigcup_{n \in \mathbb{Z}} \mathcal{F}^n \mathcal{Y}_{\mathbf{q},\infty}^{\pm}(\mathcal{R})$. From this fact combined with (4.14) and Lemma 3.3 we obtain

$$\psi_*(\tilde{\Phi}_* \mu_{\mathcal{R}}^1) = \Phi_* \nu. \quad (4.15)$$

Using the positivity of ν on all open sets, we come to the following assertion.

Lemma 4.6. *The measure $\Psi_* \mu_{\mathcal{R}}$ is positive on every open set in the phase space of the flow $(\sigma_{\mathbf{q}}, f_{\mathbf{q}})$.*

4.3 Properties of the roof function

Recall that a word $w' = w_1 \dots w_l \in \mathcal{W}_{\mathcal{A},B}$ is said to be a simple prefix of a word $w = w_1, \dots, w_l, \dots, w_n$ if $w_1 \dots w_{n-k+1} = w_k \dots w_n$ implies that either $k = 1$, or $k > l$ (see Subsection 2.1).

Lemma 4.7. *Let $\mathbf{q} \in \mathcal{W}_{\mathcal{A},B}$ be a word that has a simple positive prefix. Then the function $f_{\mathbf{q}}$ introduced in (4.13) depends only on the future, is bounded away from zero, and is Hölder continuous in the following sense: there exist positive constants $C_{\mathbf{q}}, \alpha_{\mathbf{q}}$ (depending only on $\mathbf{q}$) such that if $u = (\dots, u_{-1}, u_0, u_1 \dots) \in \mathcal{A}_{\mathbf{q}}^{\mathbb{Z}}$ and $\tilde{u} = (\dots, \tilde{u}_{-1}, \tilde{u}_0, \tilde{u}_1 \dots) \in \mathcal{A}_{\mathbf{q}}^{\mathbb{Z}}$ satisfy $u_i = \tilde{u}_i$ for $|i| \leq n$, then*

$$|f_{\mathbf{q}}(u) - f_{\mathbf{q}}(\tilde{u})| \leq C_{\mathbf{q}} \exp(-\alpha_{\mathbf{q}} n).$$

In particular, the function $f_{\mathbf{q}}$ has summable variations.

Proof. That the function $f_{\mathbf{q}}$ depends only on the future follows readily from its definition. Let us prove that it is bounded away from zero. By (4.13), (4.12) this property of $f_{\mathbf{q}}$ would follow from the same property of the function $\tau_{\mathbf{q}}^1$ defined on $\Phi^{-1}(\Omega_{\mathbf{q}})$. But from (4.8), (4.10) and Remark 4.1 one readily derives that, for $(\omega_0, \omega_1, \dots) \in \Omega_{\mathbf{q}}$,

$$\tau_{\mathbf{q}}^1(\lambda, \pi) = \log |A(\omega_0) \dots A(\omega_{s-1}) \lambda'|, \quad (\lambda, \pi) = \Phi^{-1}(\omega_0, \omega_1, \dots), \quad (4.16)$$

where s is defined in (4.9) and $(\lambda', \pi') = \mathcal{G}(\lambda, \pi) = \Phi^{-1}(\omega_s, \omega_{s+1}, \dots)$. If $\mathbf{p}$ is a simple prefix of $\mathbf{q}$, then $s \geq |\mathbf{p}|$, so that $\mathbf{p}$ is a prefix of the word

$(\omega_0, \dots, \omega_{s-1})$ and hence this word is positive. Thus all entries of the $m \times m$ matrix

$$A(\omega_0, \dots, \omega_{s-1}) = A(\omega_0) \dots A(\omega_{s-1})$$

are positive integers, while λ' is a positive vector with $|\lambda'| = 1$. Therefore, for every $(\lambda, \pi) \in \Phi^{-1}(\Omega_{\mathbf{q}})$,

$$\tau_{\mathbf{q}}^1(\lambda, \pi) = \log |A(\omega_0, \dots, \omega_{s-1})\lambda'| \geq \log m > 0.$$

We now turn to the Hölder continuity of $f_{\mathbf{q}}$. Let $u, \tilde{u}$ be as in the statement and let $\omega = \Psi_{M-B}^{-1}(u)$, $\tilde{\omega} = \Psi_{M-B}^{-1}(\tilde{u})$. By definition, $\omega, \tilde{\omega} \in \Omega_{\mathbf{q}}^{\mathbb{Z}}$, hence $(\omega_0, \omega_1, \dots), (\tilde{\omega}_0, \tilde{\omega}_1, \dots) \in \Omega_{\mathbf{q}}$. By (4.11)–(4.13)

$$f_{\mathbf{q}}(u) = \tau_{\mathbf{q}}^1(\lambda, \pi) = \log |A(\omega_0, \dots, \omega_{s-1})\lambda'|, \quad (4.17)$$

$$f_{\mathbf{q}}(\tilde{u}) = \tau_{\mathbf{q}}^1(\tilde{\lambda}, \tilde{\pi}) = \log |A(\tilde{\omega}_0, \dots, \tilde{\omega}_{s-1})\tilde{\lambda}'|, \quad (4.18)$$

where

$$\begin{aligned} (\lambda, \pi) &= \Phi^{-1}(\omega_0, \omega_1, \dots), \quad (\tilde{\lambda}, \tilde{\pi}) = \Phi^{-1}(\tilde{\omega}_0, \tilde{\omega}_1, \dots), \\ (\lambda', \pi') &= \mathcal{G}^s(\lambda, \pi), \quad (\tilde{\lambda}', \tilde{\pi}') = \mathcal{G}^s(\tilde{\lambda}, \tilde{\pi}). \end{aligned}$$

Since $\mathbf{p}$ is a simple prefix of $\mathbf{q}$, one can find $k \geq nl - 1$ such that $\omega_i = \tilde{\omega}_i$ for $i = 0, 1, \dots, k$. Then, by Remark 4.1, for some vectors $\lambda'', \tilde{\lambda}'' \in \Delta_{m-1}$,

$$\lambda = \frac{A(\omega_0, \dots, \omega_{nl-1})\lambda''}{|A(\omega_0, \dots, \omega_{nl-1})\lambda''|}, \quad \tilde{\lambda} = \frac{A(\omega_0, \dots, \omega_{nl-1})\tilde{\lambda}''}{|A(\omega_0, \dots, \omega_{nl-1})\tilde{\lambda}''|}, \quad (4.19)$$

Introduce the Hilbert metric d_H on Δ_{m-1} by

$$d_H(\lambda^{(1)}, \lambda^{(2)}) = \log \left(\max_{1 \leq i \leq m} \frac{\lambda_i^{(1)}}{\lambda_i^{(2)}} / \min_{1 \leq i \leq m} \frac{\lambda_i^{(1)}}{\lambda_i^{(2)}} \right), \quad \lambda^{(j)} = (\lambda_1^{(j)}, \dots, \lambda_m^{(j)}), j = 1, 2. \quad (4.20)$$

It is known that if an $m \times m$ non-negative matrix $A = (a_{ij})$ is such that $\sum_{j=1}^m a_{ij} > 0$ for all i , then the mapping $T_A : \Delta_{m-1} \rightarrow \Delta_{m-1}$ defined by $T_A \lambda = A\lambda/|A\lambda|$ does not increase the d_H -distance between points, while if $a_{ij} > 0$ for all i, j , then T_A is a uniform contraction (see, for example, [24]).

By definition, the word $\omega_0 \dots \omega_{nl-1}$ is a concatenation, namely, $\omega_0 \dots \omega_{nl-1} = \mathbf{p}w_1 \dots \mathbf{p}w_n$, where the word w_i , $1 \leq i \leq n$, is such that the sum of the entries in each row of the matrix $A(w_i)$ is positive. It follows (see (4.19)) that

$$d_H(\lambda, \tilde{\lambda}) \leq d_H(T_{A(\mathbf{q})}^n \lambda'', T_{A(\mathbf{q})}^n \tilde{\lambda}'') \leq C_1 \alpha^n, \quad (4.21)$$

where $C_1 \in \mathbb{R}_+$ and $\alpha \in (0, 1)$ depend only on $\mathbf{q}$. (We have used the fact that $T_{A(\mathbf{q})}$ takes Δ_{m-1} to a set of finite d_H -diameter).

Denote $A_1 = A(\omega_0, \dots, \omega_{s-1})$. Since all entries of the matrix A_1 are positive, we have (see (4.20))

$$d_H \left(\frac{A_1 \lambda}{|A_1 \lambda|}, \frac{A_1 \tilde{\lambda}}{|A_1 \tilde{\lambda}|} \right) = \log \left(\max_i \frac{(A_1 \lambda)_i}{(A_1 \tilde{\lambda})_i} / \min_i \frac{(A_1 \lambda)_i}{(A_1 \tilde{\lambda})_i} \right) \leq C_2 d_H(\lambda, \tilde{\lambda}),$$

where $(A_1 \lambda)_i$ (resp., $(A_1 \tilde{\lambda})_i$) is the i th entry of the vector $A_1 \lambda$ (resp., $A_1 \tilde{\lambda}$) and C_2 is determined by $\mathbf{q}$. Hence, by (4.17) and (4.21),

$$|f_{\mathbf{q}}(u) - f_{\mathbf{q}}(\tilde{u})| = |\log(|A_1 \lambda|/|A_1 \tilde{\lambda}|)| \leq C_2 d_H(\lambda, \tilde{\lambda}) \leq C_2 C_1 \alpha^n,$$

so it remains to set $C_{\mathbf{q}} = C_2 C_1$ and $\alpha_{\mathbf{q}} = -\log \alpha$. $\square$

4.4 Transition probabilities and the uniform expansion property.

By Theorem 3.1, the map $\mathcal{G}$ on $\Delta(\mathcal{R})$ preserves an absolutely continuous ergodic probability measure, which was denoted by ν .

Consider a word $w = w_1 \dots w_k \in \mathcal{W}_{\mathcal{A}, B}$, where $w_i = (c_i, n_i, \pi_i) \in \mathcal{A}$, $1 \leq i \leq k$. We say that w is *compatible* with a point $(\lambda, \pi) \in \Delta^\infty(\mathcal{R})$ (or (λ, π) is compatible with w) if

$$\text{either } \lambda \in \Delta_\pi^-, c_k = a, a^{n_k} \pi_k = \pi, \text{ or } \lambda \in \Delta_\pi^+, c_k = b, b^{n_k} \pi_k = \pi.$$

Assuming that w is compatible with (λ, π) , we set

$$t_w(\lambda, \pi) = \left(\frac{A(w)\lambda}{|A(w)\lambda|}, \pi_1 \right). \quad (4.22)$$

From the definition of $\mathcal{G}$ (see (3.6)) it follows that

$$\mathcal{G}^{-n}(\lambda, \pi) = \{t_w(\lambda, \pi) : |w| \text{ and } w \text{ is compatible with } (\lambda, \pi)\}. \quad (4.23)$$

Note that the set $\mathcal{G}^{-n}(\lambda, \pi)$ is infinite.

In Subsection 4.1 we introduced, for each word $w \in \mathcal{W}_{\mathcal{A}, B}$, the set $\Delta(w) \subset \Delta^\infty(\mathcal{R})$ (see (4.5)). One can readily check (see Remark 4.1) that

$$\Delta(w) = \{t_w(\lambda, \pi) : (\lambda, \pi) \in \Delta^\infty(\mathcal{R}) \text{ is compatible with } w\}. \quad (4.24)$$

For every $n \in \mathbb{N}$, we have the ν -measurable partition $\mathcal{G}^{-n}\varepsilon$ of $\Delta^\infty(\mathcal{R})$, where ε is the partition into separate points. Each element of $\mathcal{G}^{-n}\varepsilon$ is $\mathcal{G}^{-n}(\lambda, \pi)$ for some $(\lambda, \pi) \in \Delta^\infty(\mathcal{R})$, its points correspond to the words $w \in \mathcal{W}_{\mathcal{A},B}$ of length n compatible with (λ, π) and have the form $t_w(\lambda, \pi)$ (see (4.22), (4.23)). We denote by $\nu(w|(\lambda, \pi))$ the conditional measure (determined by ν) of the point corresponding to w , given the element $\mathcal{G}^{-n}(\lambda, \pi)$ of the partition $\mathcal{G}^{-n}\varepsilon$.

In Section 3.5 of [4] it is proved that if w is compatible with (λ, π) , then

$$\nu(w|(\lambda, \pi)) = \frac{\rho(t_w(\lambda, \pi))}{\rho(\lambda, \pi)|A(w)\lambda|^m}. \quad (4.25)$$

Now consider the set $\Delta(\mathbf{q})$ corresponding to a word $\mathbf{q} \in \mathcal{W}_{\mathcal{A},B}$ (see (4.5)). Every point from $\Delta(\mathbf{q})$ is of the form $(\lambda, \pi_{\mathbf{q}})$, where $\pi_{\mathbf{q}}$ is a fixed permutation and λ belongs to a set $\Delta'(\mathbf{q}) \subset \Delta_{m-1}$. Denote by $d(\mathbf{q})$ the diameter of $\Delta'(\mathbf{q})$ with respect to the Hilbert metric on Δ_{m-1} introduced in (4.20).

Proposition 4.8. *There are positive constants β_1 and β_2 (depending only on $\mathcal{R}$) such that for every positive word $\mathbf{p}' \in \mathcal{W}_{\mathcal{A},B}$ with $d(\mathbf{p}') \leq \beta_1$, the following holds. Let $\mathbf{p}$ be a word that has $\mathbf{p}'$ as a simple prefix, and let a word $\mathbf{r} \in \mathcal{W}_{\mathcal{A},B}$ start and end with $\mathbf{p}$ and contain no other occurrences of $\mathbf{p}$. Then for any $(\lambda, \pi) \in \Delta(\mathbf{r}) \cap \Phi^{-1}(\Omega_{\mathbf{p}})$, we have*

$$\left| \frac{\nu(\Delta(\mathbf{r})) \exp[m\tau_{\mathbf{p}}^1(\lambda, \pi)]}{\nu(\Delta(\mathbf{p}))} - 1 \right| \leq \beta_2 d(\mathbf{p}'),$$

where $\tau_{\mathbf{p}}^1$ is defined by (4.16).

Proof. By assumption, the word $\mathbf{r}$ has the form $\mathbf{r} = \mathbf{p}'u\mathbf{p} = \mathbf{p}\tilde{u}$ for some $u, \tilde{u} \in \mathcal{W}_{\mathcal{A},B}$. Hence

$$\nu(\Delta(\mathbf{r})) = \int_{\Delta(\mathbf{p})} \nu(\mathbf{p}'u|(\lambda, \pi)) d\nu(\lambda, \pi). \quad (4.26)$$

From (4.25) we obtain

$$\nu(\mathbf{p}'u|(\lambda, \pi)) = \frac{\rho(t_{\mathbf{p}'u}(\lambda, \pi))}{\rho(\lambda, \pi)} \cdot \frac{1}{|A(\mathbf{p}'u)\lambda|^m}. \quad (4.27)$$

By (4.20), taking into account that $|\lambda| = 1$ when $\lambda \in \Delta_{m-1}$, for any two points, $(\lambda, \pi_{\mathbf{p}'})$, $(\tilde{\lambda}, \pi_{\mathbf{p}'}) \in \Delta(\mathbf{p}')$, and for $i = 1, \dots, m$, we have

$$e^{-d(\mathbf{p}')} \leq \tilde{\lambda}_i / \lambda_i \leq e^{d(\mathbf{p}')}.$$

Let us estimate the first ratio on the right-hand side of (4.27). From the fact that $\mathbf{p}'u\mathbf{p}' \in \mathcal{W}_{\mathcal{A},B}$ it follows that if $(\lambda, \pi_{\mathbf{p}'}) \in \Delta(\mathbf{p}')$, then $(\lambda, \pi_{\mathbf{p}'})$ is compatible with $\mathbf{p}'u$. Hence $t_{\mathbf{p}'u} \in \Delta(\mathbf{p}'u) \subset \Delta(\mathbf{p}')$ (see (4.24)). Denote

$$(\hat{\lambda}, \hat{\pi}) := t_{\mathbf{p}'u}(\lambda, \pi_{\mathbf{p}'}), \quad \hat{\lambda} = (\hat{\lambda}_1, \dots, \hat{\lambda}_m),$$

and observe that $\hat{\pi} = \pi_{\mathbf{p}'}$. By (4.28)

$$e^{-d(\mathbf{p}')} \lambda_i \leq \hat{\lambda}_i \leq e^{d(\mathbf{p}')} \lambda_i, \quad i = 1, \dots, m,$$

while by Theorem 3.1 $\rho(\hat{\lambda}, \hat{\pi}) = P_{\hat{\pi}}(\hat{\lambda})/Q_{\hat{\pi}}(\hat{\lambda})$, where $P_{\hat{\pi}}$ and $Q_{\hat{\pi}}$ are homogeneous polynomials with non-negative coefficients. Therefore,

$$P_{\hat{\pi}}(\hat{\lambda}) \leq P_{\hat{\pi}}(e^{d(\mathbf{p}')} \lambda) \leq e^{\gamma_1 d(\mathbf{p}')} P_{\hat{\pi}}(\lambda),$$

$$Q_{\hat{\pi}}(\hat{\lambda}) \geq Q_{\hat{\pi}}(e^{-d(\mathbf{p}')} \lambda) \geq e^{-\gamma_2 d(\mathbf{p}')} Q_{\hat{\pi}}(\lambda),$$

where γ_1 and γ_2 are determined by ρ (and, eventually, by $\mathcal{R}$). From this we immediately obtain

$$\exp[-(\gamma_1 + \gamma_2)d(\mathbf{p}')] \leq \frac{\rho(t_{\mathbf{p}'u}(\lambda, \pi))}{\rho(\lambda, \pi)} \leq \exp[(\gamma_1 + \gamma_2)d(\mathbf{p}')] \quad (4.29)$$

Using (4.28) and the positivity of all entries of the matrix $A(\mathbf{p}'u)$, for any $(\lambda, \pi_{\mathbf{p}'})$, $(\tilde{\lambda}, \pi_{\mathbf{p}'}) \in \Delta(\mathbf{p}')$, we have

$$\exp[-d(\mathbf{p}')] \leq \frac{|A(\mathbf{p}'u)\tilde{\lambda}|}{|A(\mathbf{p}'u)\lambda|} \leq \exp[d(\mathbf{p}')] \quad (4.30)$$

Now fix an arbitrary point $(\lambda^0, \pi^0) \in \Delta(\mathbf{r}) \cap \Phi^{-1}(\Omega_{\mathbf{p}})$. By Lemma 4.2 $(\lambda^0, \pi^0) = \Phi^{-1}(\omega_0, \omega_1, \dots)$ for some $(\omega_0, \omega_1, \dots) \in \Omega_{\mathbf{p}}$. Let s be defined by (4.9) and $(\lambda', \pi') := \mathcal{G}^s(\lambda^0, \pi^0)$. Clearly, $s = |\mathbf{p}'u|$, hence $(\lambda', \pi') \in \Delta(\mathbf{p}) \subset \Delta(\mathbf{p}')$, and by (4.26), (4.27), (4.29), (4.30),

$$\begin{aligned} \frac{\nu(\Delta(\mathbf{r}))|A(\mathbf{p}'u)\lambda'|^m}{\nu(\Delta(\mathbf{p}))} &= \frac{1}{\nu(\Delta(\mathbf{p}))} \int_{\Delta(\mathbf{p})} \frac{\rho(t_{\mathbf{p}'u}(\lambda, \pi))}{\rho(\lambda, \pi)} \cdot \frac{|A(\mathbf{p}'u)\lambda'|^m}{|A(\mathbf{p}'u)\lambda|^m} d\nu(\lambda, \pi) \\ &\leq \exp[(\gamma_1 + \gamma_2 + 1)d(\mathbf{p}')] \end{aligned} \quad (4.31)$$

Similarly,

$$\frac{\nu(\Delta(\mathbf{r}))|A(\mathbf{p}'u)\lambda'|^m}{\nu(\Delta(\mathbf{p}))} \geq \exp[-(\gamma_1 + \gamma_2 + 1)d(\mathbf{p}')] \quad (4.32)$$

From (4.31), (4.32) we obtain

$$\left| \frac{\nu(\Delta(\mathbf{r}))|A(\mathbf{p}'u)\lambda'|^m}{\nu(\Delta(\mathbf{p}))} - 1 \right| \leq \exp[(\gamma_1 + \gamma_2 + 1)d(\mathbf{p})] \leq 2(\gamma_1 + \gamma_2 + 1)d(\mathbf{p}'),$$

where the last inequality holds when $(\gamma_1 + \gamma_2 + 1)d(\mathbf{p}') \leq \log 2$. It remains to recall that by (4.16) $|A(\mathbf{p}'u)\lambda'| = \exp(\tau_{\mathbf{p}}^1(\lambda, \pi))$. $\square$

5 Zippered rectangles and Abelian differentials. Completion of the proof of Theorem 1.1

Fix a connected component $\mathcal{H}$ of the space $\mathcal{M}_\kappa$ (see Section 1). To this component there corresponds a unique Rauzy class $\mathcal{R}$ in such a way that the following is true [21, 13].

Theorem 5.1 (Veech). *There exists a finite-to-one measurable map $\pi_{\mathcal{R}} : \tilde{\mathcal{V}}(\mathcal{R}) \rightarrow \mathcal{H}$ such that $\pi_{\mathcal{R}} \circ P^t = g_t \circ \pi_{\mathcal{R}}$ and $(\pi_{\mathcal{R}})_*\mu_{\mathcal{R}} = \mu_\kappa$ for all $t \in \mathbb{R}$.*

(Recall that the set $\tilde{\mathcal{V}}(\mathcal{R})$ is defined in Subsection 3.2).

Corollary 5.2. 1. *If η is a $\{g_t\}$ -invariant ergodic probability measure on $\mathcal{H}$, then there exists a $\{P^t\}$ -invariant measure $\tilde{\eta}$ on $\tilde{\mathcal{V}}(\mathcal{R})$ with $(\pi_{\mathcal{R}})_*\tilde{\eta} = \eta$.*

2. *If $\tilde{\eta}$ is a $\{P^t\}$ -invariant probability measure on $\tilde{\mathcal{V}}(\mathcal{R})$ such that the $\{g_t\}$ -invariant measure $(\pi_{\mathcal{R}})_*\tilde{\eta}$ is ergodic, then*

$$h_{\tilde{\eta}}(\{P^t\}) = h_{(\pi_{\mathcal{R}})_*\tilde{\eta}}(\{g_t\}). \quad (5.1)$$

Proof. 1. Let η be an ergodic $\{g_t\}$ -invariant probability measure on $\mathcal{H}$. By ergodicity, there is a set $\mathcal{H}' \subset \mathcal{H}$ such that $\eta(\mathcal{H}') = 1$ and the cardinality of the preimage $\pi_{\mathcal{R}}^{-1}(p)$ does not depend on $p \in \mathcal{H}'$. The sets $\pi_{\mathcal{R}}^{-1}(p)$ form a measurable partition of $\tilde{\mathcal{V}}(\mathcal{R})$. By assigning equal weights (conditional measures) to all points in $\pi_{\mathcal{R}}^{-1}(p)$, $p \in \mathcal{H}$, we obtain a $\{P^t\}$ -invariant probability measure $\tilde{\eta}$ on $\tilde{\mathcal{V}}(\mathcal{R})$ such that $(\pi_{\mathcal{R}})_*\tilde{\eta} = \eta$.

2. Let $\tilde{\eta}$ be a $\{P^t\}$ -invariant probability measure on $\tilde{\mathcal{V}}(\mathcal{R})$ and $\eta = (\pi_{\mathcal{R}})_*\tilde{\eta}$. Assume that η is ergodic with respect to the flow $\{g_t\}$, hence it is ergodic with respect to the automorphism g_{t_0} for some $t_0 > 0$.

Denote by $\mathfrak{c}$ the canonical partition for P^{t_0} and $\tilde{\eta}$, i.e., the partition of $\tilde{\mathcal{V}}(\mathcal{R})$ corresponding to the decomposition of P^{t_0} into ergodic components

with respect to $\tilde{\eta}$. For an element C of $\mathfrak{c}$, denote by $\tilde{\eta}_C$ the conditional measure induced by $\tilde{\eta}$ on C , and consider $\tilde{\eta}_C$ as a measure on $\tilde{\mathcal{V}}(\mathcal{R})$. Let us note that for $\tilde{\eta}$ -almost all C , the measure $\tilde{\eta}_C$ exists and is ergodic with respect to P^{t_0} .

Take a measurable set M comprised of preimages $\pi_{\mathcal{R}}^{-1}(p)$. From the Birkhoff ergodic theorem, applied to P^{t_0} , $\tilde{\eta}$, M and to P^{t_0} , $\tilde{\eta}_C$, M , it follows that $\mathfrak{c}$ and the partition of $\tilde{\mathcal{V}}(\mathcal{R})$ into the preimages $\pi_{\mathcal{R}}^{-1}(p)$, $p \in \mathcal{H}$, are independent with respect to the measure $\tilde{\eta}$. Therefore $(\pi_{\mathcal{R}})_*\tilde{\eta}_C = \eta$ for $\tilde{\eta}$ -almost all elements $C \in \mathfrak{c}$.

As before, the cardinality of $\pi_{\mathcal{R}}^{-1}(p)$ is a constant, say k , on a set $\mathcal{H}' \subset \mathcal{H}$ with $\eta(\mathcal{H}') = 1$. Moreover, if $\tilde{\eta}_C$ is P^{t_0} -ergodic, then the conditional measure induced by $\tilde{\eta}_C$ on $\pi_{\mathcal{R}}^{-1}(p)$ for η -almost all $p \in \mathcal{H}'$ is uniform. Therefore, the measure space $(\mathcal{V}(\mathcal{R}), \tilde{\eta}_C)$ is isomorphic to the direct product of $(\mathcal{H}, \eta)$ and a set consisting of k points of mass $1/k$ each. In this representation, P^{t_0} becomes a skew product with base $(\mathcal{H}, \eta, g_{t_0})$, and the Abramov-Rokhlin formula [1] for the entropy of a skew product implies that $h_{\tilde{\eta}_C}(P^{t_0}) = h_{\eta}(g_{t_0})$. Thus the entropy of $\tilde{\eta}$ -almost every ergodic component of P^{t_0} equals $h_{\eta}(g_{t_0})$, which implies that $h_{\tilde{\eta}}(P^{t_0}) = h_{\eta}(g_{t_0})$. Since the entropy of automorphisms forming a measurable flow $\{S_t\}$ with respect to an $\{S_t\}$ -invariant probability measure μ satisfies the equation $h_{\mu}(\{S_t\}) = |t|h_{\mu}(S_1)$, we come to (5.1). $\square$

From Theorem 5.1 and Corollary 5.2, taking into account the ergodicity of the measure μ_{κ} with respect to $\{g_t\}$ (see Section 1), we readily obtain the following.

Corollary 5.3. *To prove that μ_{κ} is a unique measure with maximal entropy for the flow $\{g_t\}$ it suffices to show that $\mu_{\mathcal{R}}$ is a unique measure with maximal entropy for the flow $\{P^t\}$.*

Recall that the Rauzy class $\mathcal{R}$ we deal with consists of permutations on m symbols. With this in mind we derive from (1.1) and Corollary 5.2 that

$$h_{\mu_{\mathcal{R}}}(\{P^t\}) = 2g - 1 + r = m, \quad (5.2)$$

We call a point $x \in \tilde{\mathcal{V}}(\mathcal{R})$ *infinitely renormalizable* if its trajectory $\{P^t x, t \in \mathbb{R}\}$ intersects the transversal $\mathcal{Y}^{\pm}(\mathcal{R})$ infinitely many times both for $t > 0$ and for $t < 0$. The set of infinitely renormalizable points was denoted in Section 3.2 by $\mathcal{Y}_{\infty}^{\pm}(\mathcal{R})$.

The following proposition is in essence contained in [21] and [15].

Proposition 5.4. *There exists a Borel measurable set $V \subset \mathcal{H}$ such that*

- (i) $\mu(V) = 1$ for every ergodic $\{g_t\}$ -invariant probability measure μ on $\mathcal{H}$;
- (ii) $\pi_{\mathcal{R}}^{-1}(p) \cap \mathcal{Y}_{\infty}^{\pm} \neq \emptyset$ for each $p \in V$.

Proof. For a compact set $K \subset \mathcal{H}$, denote by $K^{\pm}$ the set of points $p \in \mathcal{H}$ for which there exist $t_n \rightarrow +\infty$ and $s_n \rightarrow -\infty$ such that $g_{t_n}p \in K$, $g_{s_n}p \in K$ for $n = 1, 2, \dots$.

Take an increasing sequence of compact sets K_n such that $\bigcup_n K_n = \mathcal{H}$, and let $V = \bigcup_n K_n^{\pm}$. The set V is obviously Borel measurable. By definition, for every probability measure μ on $\mathcal{H}$, there exists n_0 with $\mu(K_{n_0}) > 0$. If, in addition, μ is $\{g_t\}$ -invariant and ergodic, then $\mu(K_{n_0}^{\pm}) = 1$ and hence $\mu(V) = 1$.

Let (σ, ω) belong to an equivalence class $p \in V$ (see Section 1). Then, by Masur's theorem [15], the foliations corresponding to $\Re(\omega)$ and $\Im(\omega)$ are both uniquely ergodic. This implies, in particular, the existence of an infinitely renormalizable zippered rectangle in $\pi_{\mathcal{R}}^{-1}p$ (see, for instance, [27]), which is all that we had to prove. $\square$

Lemma 5.5. *Let $\tilde{\eta}$ be an ergodic $\{P^t\}$ -invariant probability measure on $\mathcal{V}(\mathcal{R})$. Then there exists a positive word $\mathbf{p} \in \mathcal{W}_{\mathcal{A}, B}$ such that $\tilde{\eta}(\mathcal{V}_{\mathbf{p}}(\mathcal{R})) = 1$. Moreover, if $h_{\tilde{\eta}}(\{P^t\}) > 0$, then for any $\mathbf{p}$ such that $\tilde{\eta}(\mathcal{V}_{\mathbf{p}}(\mathcal{R})) = 1$, there also exists a word $\mathbf{q} \in \mathcal{W}_{\mathcal{A}, B}$ such that $\tilde{\eta}(\mathcal{V}_{\mathbf{q}}(\mathcal{R})) = 1$ and $\mathbf{p}$ is a simple prefix of $\mathbf{q}$.*

Proof. Since the measure $\tilde{\eta}$ is ergodic with respect to $\{P^t\}$, its projection $\eta := (\pi_{\mathcal{R}})_* \tilde{\eta}$ is ergodic with respect to $\{g_t\}$. Let V be as in Proposition 5.4 and $W = \pi_{\mathcal{R}}^{-1}(V)$. By Proposition 5.4 $\eta(V) = 1$, which implies that $\tilde{\eta}(W) = 1$.

For every point $x = (\lambda, \pi, \delta) \in W$, the interval exchange (λ, π) is uniquely ergodic. Now Lemma 4.3 yields a positive word $\mathbf{q}_x \in \mathcal{W}_{\mathcal{A}, B}$ such that $(\lambda, \pi) \in \Delta(\mathbf{q}_x)$. Since $\eta(W) = 1$, while $\mathcal{W}_{\mathcal{A}, B}$ is countable, there exists $\mathbf{q} \in \mathcal{W}_{\mathcal{A}, B}$ such that $\tilde{\eta}(\{x : \mathbf{q}_x = \mathbf{q}\}) > 0$. For this $\mathbf{q}$ we have $\tilde{\eta}(\mathcal{Y}_{\mathbf{q}, \infty}^{\pm})(\mathcal{R}) > 0$ and, since $\tilde{\eta}$ is ergodic, $\tilde{\eta}(\mathcal{V}_{\mathbf{q}}(\mathcal{R})) = 1$.

Let us now consider the set $\mathcal{W}(\mathbf{p}) \subset \mathcal{W}_{\mathcal{A}, B}$ of all words that have $\mathbf{p}$ as a prefix. If $\mathbf{q} \in \mathcal{W}(\mathbf{p})$, and $\mathbf{p}$ is not a simple prefix of $\mathbf{q}$, then $\mathbf{q}$ is a concatenation: $\mathbf{q} = \mathbf{q}'\mathbf{q}'' \dots \mathbf{q}'\mathbf{q}''$, where $\mathbf{q}'$ is a prefix of $\mathbf{p}$, and $\mathbf{q}''$ is either a prefix of $\mathbf{q}'$ or empty. In this situation, either $\tilde{\eta}(\mathcal{V}_{\mathbf{q}}(\mathcal{R})) = 0$ or $\tilde{\eta}(\mathcal{V}_{\mathbf{q}}(\mathcal{R})) > 0$ and $\tilde{\eta}$ is concentrated on periodic points of the flow $\{P^t\}$. But this can not be the case for all $\mathbf{q} \in \mathcal{W}(\mathbf{p})$, since $\tilde{\eta}(\mathcal{V}_{\mathbf{q}}(\mathcal{R})) = 1$ and $h_{\tilde{\eta}}(\{P^t\}) > 0$. $\square$

The following statement will be also used below.

Lemma 5.6. *Let $\Gamma = (V, E)$ be a directed graph with $|V| = \infty$ and $W(\Gamma)$ be the corresponding family of words (see Subsection 2.1). Then for each $w \in W(\Gamma)$ and each $n \in \mathbb{N}$, there exists a word $w' \in W(\Gamma)$ of the form $w' = ww_1ww_2w, \dots, w_{n-1}w$, $w_i \in W(\Gamma)$, containing n disjoint subwords equal to w and such that the word $w'' := ww_1ww_2w, \dots, w_{n-1}$ is simple.*

Proof. Denote the first and last letters of w by v^- and v^+ respectively, and construct by induction a sequence of letters $v_1, v_2, \dots$ as follows. For v_1 we take an arbitrary letter that is not contained in w . If $v_1, \dots, v_k$, $1 \leq k \leq n-1$, are already chosen, we denote by w_k^+ (respectively, w_k^-) the shortest word from v^+ to v_k (resp., from v_k to v^-). (If there are several words of the same, minimal, length, we take any of them.) Then take for v_{k+1} an arbitrary vertex except for those contained in at least one of the words $w, w_1^+, w_1^-, \dots, w_k^+, w_k^-$.

Consider the sequence of words

$$w, w_1^+, w_1^-, w, w_2^+, w_2^-, w, \dots, w, w_n^+, w_{n-1}^-, w.$$

Delete the last letter from each of these words but the last w , and denote by w' the concatenation of the words thus obtained. It is easy to check that w' possesses the required properties. (To define w_k one should remove the first and last letters from w_k^+ , the last letter from w_k^- , and then take the concatenation of the two words obtained.) $\square$

End of the proof of Theorem 1.1. By Corollary 5.3, it suffices to prove that $\mu_{\mathcal{R}}$ is a unique measure with maximal entropy for the flow $\{P^t\}$.

Let μ be a measure on $\tilde{V}(\mathcal{R})$ with $h_\mu(\{P^t\}) \geq h_{\mu_{\mathcal{R}}}(\{P^t\})$. Without loss of generality we can assume that μ is ergodic (otherwise one could pass to an ergodic component). By Lemma 5.5, there exists a word $\mathbf{q} \in \mathcal{W}_{\mathcal{A}, \mathcal{B}}$ that has a simple positive prefix and is such that $\mu(\mathcal{V}_{\mathbf{q}}(\mathcal{R})) = 1$. Remind that $\mu_{\mathcal{R}}(\mathcal{V}_{\mathbf{q}}(\mathcal{R})) = 1$ as well. By Lemma 4.5, the flow $\{P^t\}|_{\mathcal{V}_{\mathbf{q}}(\mathcal{R})}$ is embedded in the suspension flow $(\sigma_{\mathbf{q}}, f_{\mathbf{q}})$ via a mapping Ψ . The roof function $f_{\mathbf{q}}$ clearly depends only on the future (see Subsection 2.1). Moreover, by Lemma 4.7 $f_{\mathbf{q}}$ is bounded away from zero and has summable variations.

By Lemma 4.6 the measure $\Psi_*\mu_{\mathcal{R}}$ is positive on all non-empty open subsets of the phase space of the flow $(\sigma_{\mathbf{q}}, f_{\mathbf{q}})$. This measure induces, in a canonical way, a probability measure $\bar{\mu}_{\mathbf{q}}$ on $(\mathcal{A}_{\mathbf{q}})^{\mathbb{Z}}$, the base of the suspension flow $(\sigma_{\mathbf{q}}, f_{\mathbf{q}})$. It follows that $\bar{\mu}_{\mathbf{q}}$ is positive on all cylinders in $(\mathcal{A}_{\mathbf{q}})^{\mathbb{Z}}$.

Let us prove that $\bar{\mu}_{\mathbf{q}}$ satisfies the other conditions imposed on the measure $\bar{\mu}$ in Theorem 2.2. (We apply this theorem to the complete graph with vertex set $\mathcal{A}_{\mathbf{q}}$; in this situation every sequence of letters (i.e., vertices of the graph) is a word.)

Apply Lemma 5.6 to the connected graph $\Gamma_{\mathcal{A},B}$ (see Subsection 4.1) and to the word $\mathbf{q}$ taken as w . From this lemma we obtain, for each $n \in \mathbb{N}$, a word of the form $\mathbf{q}q_1\mathbf{q} \dots \mathbf{q}q_n\mathbf{q} \in \mathcal{W}_{\mathcal{A},B}$ whose prefix $\mathbf{q}q_1\mathbf{q} \dots \mathbf{q}q_n$ is a simple word.

Each word $\mathbf{a}_i := \mathbf{q}q_i\mathbf{q}$, $i = 1, \dots, n$, is a letter in the alphabet $\mathcal{A}_{\mathbf{q}}$ introduced in Subsection 4.2, and $\mathbf{a} := (\mathbf{a}_1, \dots, \mathbf{a}_n)$ is clearly a simple word for all n . Denote by $\hat{a}$ an arbitrary word in the alphabet $\mathcal{A}_{\mathbf{q}}$ that does not contain $\mathbf{a}$ as a subword, and consider the cylinders $C_{\mathbf{a}}$ and $C_{\mathbf{a}\hat{a}\mathbf{a}} \subset (\mathcal{A}_{\mathbf{q}})^{\mathbb{Z}}$. Using the definition of the measures and maps that appear below, we have

$$\frac{\bar{\mu}_{\mathbf{q}}(C_{\mathbf{a}\hat{a}\mathbf{a}})}{\bar{\mu}_{\mathbf{q}}(C_{\mathbf{a}})} = \frac{(\tilde{\Phi}_*\mu_{\mathcal{R},\mathbf{q}}^1)(\Psi_{M-B}^{-1}C_{\mathbf{a}\hat{a}\mathbf{a}})}{(\tilde{\Phi}_*\mu_{\mathcal{R},\mathbf{q}}^1)(\Psi_{M-B}^{-1}C_{\mathbf{a}})} = \frac{(\tilde{\Phi}_*\mu_{\mathcal{R}}^1)(\Psi_{M-B}^{-1}C_{\mathbf{a}\hat{a}\mathbf{a}})}{(\tilde{\Phi}_*\mu_{\mathcal{R}}^1)(\Psi_{M-B}^{-1}C_{\mathbf{a}})}. \quad (5.3)$$

Observe that $\Psi_{M-B}^{-1}C_{\mathbf{a}} = C \cap \Omega_{\mathbf{q}}^{\mathbb{Z}}$, where C is a cylinder in $\Omega_{\mathcal{A},B}^{\mathbb{Z}}$ whose support belongs to $\mathbb{Z}_+$. It follows that

$$\psi^{-1}(\psi(C \cap \Omega_{\mathbf{q}}^{\mathbb{Z}})) = C \cap \Omega_{\mathbf{q}}^{\mathbb{Z}_+},$$

where $\Omega_{\mathbf{q}}^{\mathbb{Z}_+}$ consists of all $\omega = (\omega_k, k \in \mathbb{Z}) \in \Omega_{\mathcal{A},B}^{\mathbb{Z}}$ such that $(\omega_n, \dots, \omega_{n+|\mathbf{q}|-1}) = \mathbf{q}$ for $n = 0$ and for infinitely many $n > 0$. Hence

$$\begin{aligned} (\psi_*(\tilde{\Phi}_*\mu_{\mathcal{R}}^1))(\psi\Psi_{M-B}^{-1}C_{\mathbf{a}}) &= (\psi_*(\tilde{\Phi}_*\mu_{\mathcal{R}}^1))(\psi(C \cap \Omega_{\mathbf{q}}^{\mathbb{Z}})) \\ &= (\tilde{\Phi}_*\mu_{\mathcal{R}}^1)(C \cap \Omega_{\mathbf{q}}^{\mathbb{Z}_+}). \end{aligned}$$

Since the measure $\tilde{\Phi}_*\mu_{\mathcal{R}}^1$ is shift-invariant, while the word $\mathbf{q}$ is a prefix of $\mathbf{a}_1 \in \mathcal{W}_{\mathcal{A},B}$, from the Poincaré Recurrence Theorem we obtain

$$(\tilde{\Phi}_*\mu_{\mathcal{R}}^1)(C \cap \Omega_{\mathbf{q}}^{\mathbb{Z}}) = (\tilde{\Phi}_*\mu_{\mathcal{R}}^1)(C \cap \Omega_{\mathbf{q}}^{\mathbb{Z}_+}) = (\tilde{\Phi}_*\mu_{\mathcal{R}}^1)(C),$$

so that

$$(\psi_*(\tilde{\Phi}_*\mu_{\mathcal{R}}^1))(\psi\Psi_{M-B}^{-1}C_{\mathbf{a}}) = (\tilde{\Phi}_*\mu_{\mathcal{R}}^1)(\Psi_{M-B}^{-1}C_{\mathbf{a}}). \quad (5.4)$$

In this argument one may replace $C_{\mathbf{a}}$ by $C_{\mathbf{a}\hat{a}\mathbf{a}}$ to obtain

$$(\psi_*(\tilde{\Phi}_*\mu_{\mathcal{R}}^1))(\psi\Psi_{M-B}^{-1}C_{\mathbf{a}\hat{a}\mathbf{a}}) = (\tilde{\Phi}_*\mu_{\mathcal{R}}^1)(\Psi_{M-B}^{-1}C_{\mathbf{a}\hat{a}\mathbf{a}}). \quad (5.5)$$

Substitution of (5.4) and (5.5) in (5.3) with taking into account (4.15) yields

$$\frac{\bar{\mu}_{\mathbf{q}}(C_{\mathbf{a}\hat{a}\mathbf{a}})}{\bar{\mu}_{\mathbf{q}}(C_{\mathbf{a}})} = \frac{(\Phi_*\nu)(\psi\Psi_{M-B}^{-1}C_{\mathbf{a}\hat{a}\mathbf{a}})}{(\tilde{\Phi}_*\nu)(\psi\Psi_{M-B}^{-1}C_{\mathbf{a}})} = \frac{\nu(\Phi^{-1}(\psi\Psi_{M-B}^{-1}C_{\mathbf{a}\hat{a}\mathbf{a}}))}{\nu(\Phi^{-1}(\psi\Psi_{M-B}^{-1}C_{\mathbf{a}}))}. \quad (5.6)$$

In a similar way as above we have

$$\psi(\Psi_{M-B}^{-1}C_{\mathbf{a}}) = C_{\mathbf{a}(\mathbf{q})} \cap \Omega_{\mathbf{q}}, \quad \psi(\Psi_{M-B}^{-1}C_{\mathbf{a}\hat{a}\mathbf{a}}) = C_{\hat{\mathbf{a}}(\mathbf{q})} \cap \Omega_{\mathbf{q}},$$

where $C_{\mathbf{a}(\mathbf{q})}$, $C_{\hat{\mathbf{a}}(\mathbf{q})}$ are the cylinders in $\Omega_{\mathcal{A},B}$ corresponding to the words $\mathbf{a}(\mathbf{q}) := \mathbf{q}q_1\mathbf{q} \dots \mathbf{q}q_n\mathbf{q} \in \mathcal{W}_{\mathcal{A},B}$ and $\hat{\mathbf{a}}(\mathbf{q}) := \mathbf{a}(\mathbf{q})\hat{w}\mathbf{a}(\mathbf{q}) \in \mathcal{W}_{\mathcal{A},B}$ with some $\hat{w} \in \Omega_{\mathcal{A},B}$, respectively. Therefore (see (5.6), (4.5)),

$$\frac{\bar{\mu}_{\mathbf{q}}(C_{\mathbf{a}\hat{a}\mathbf{a}})}{\bar{\mu}_{\mathbf{q}}(C_{\mathbf{a}})} = \frac{\nu(\Delta(\hat{\mathbf{a}}(\mathbf{q})))}{\nu(\Delta(\mathbf{a}(\mathbf{q})))}. \quad (5.7)$$

Let us now apply Proposition 4.8 with $\mathbf{p}' = \mathbf{q}q_1\mathbf{q} \dots \mathbf{q}q_n\mathbf{q}$, $\mathbf{p} = \mathbf{a}(\mathbf{q})$, $\mathbf{r} = \hat{\mathbf{a}}(\mathbf{q})$, where n is large enough. We may do so because of the following two facts: 1) the choice of the word $\hat{a}$ above implies that $\hat{w}$ does not contain subwords equal to $\mathbf{a}(\mathbf{q})$; 2) $d(\mathbf{p}) \leq e^{-n\alpha}$ for some $\alpha > 0$, as can be shown in the manner of Subsection 4.3. By this proposition combined with (5.3) we obtain

$$\left| \frac{\bar{\mu}_{\mathbf{q}}(C_{\mathbf{a}\hat{a}\mathbf{a}})}{\bar{\mu}_{\mathbf{q}}(C_{\mathbf{a}})} - \exp(-m\tau_{\mathbf{a}(\mathbf{q})}^1(\lambda, \pi)) \right| \leq \beta_2 \exp(-n\alpha - m\tau_{\mathbf{q}}^1(\lambda, \pi)),$$

where

$$(\lambda, \pi) \in (\Phi^{-1}\psi\Psi_{M-B}C_{\mathbf{a}\hat{a}\mathbf{a}}) \cap \Phi^{-1}\Omega_{\mathbf{a}(\mathbf{q})}.$$

Using (2.3), (4.13), and (4.10)–(4.12), one can check that if $u \in C_{\mathbf{a}\hat{a}\mathbf{a}}$ is such that $(u_n, \dots, u_{n+|\mathbf{a}|-1}) = \mathbf{a}$ for infinitely many $n > 0$ and $n < 0$, then $\tilde{\tau}(u, C_{\mathbf{a}}) = \tau_{\mathbf{a}(\mathbf{q})}(\lambda, \pi)$, where $(\lambda, \pi) = \Phi^{-1}\psi\Psi_{M-B}u \in \Phi^{-1}\Omega_{\mathbf{a}(\mathbf{q})}$. It remains to note that by the Recurrence Theorem, $\bar{\mu}_{\mathbf{q}}$ -almost all $u \in C_{\mathbf{a}\hat{a}\mathbf{a}}$ satisfy the stated condition.

We thus see that the measure $\bar{\mu}_{\mathbf{q}}$ satisfies the assumptions of Theorem 2.2 (in particular, the constant s in (2.4) equals $m = h_{\Psi_*\mu_{\mathcal{R}}}(\sigma_{\mathbf{q}}, f_{\mathbf{q}})$). This theorem now implies that $(\bar{\mu}_{\mathbf{q}})_f$ is a measure with maximal entropy for $(\sigma_{\mathbf{q}}, f_{\mathbf{q}})$ and hence $\mu_{\mathcal{R}}$ is a measure with maximal entropy for $\{P^t\}$, so that $h_{\text{top}}(\{P^t\}) = h_{\text{top}}(\sigma_{\mathbf{q}}, f_{\mathbf{q}}) = m$.

As to the measure μ , it follows that $h_{\mu}(\{P^t\}) = h_{\mu_{\mathcal{R}}}(\{P^t\})$. Then $\Psi_*\mu$ has the same entropy with respect to the suspension flow $(\sigma_{\mathbf{q}}, f_{\mathbf{q}})$ (see (5.2)) and

hence is a measure with maximal entropy for this flow. But from Theorem 2.2 we know that such a measure is unique. This completes the proof of Theorem 1.1.

Acknowledgements. A.I.B. is supported in part by Grant MK-4893.2010.1 of the President of the Russian Federation, by the Programme on Mathematical Control Theory of the Presidium of the Russian Academy of Sciences, by the Russian Foundation for Basic Research under Grant 10-01-00739-, by the Russian Ministry of Education and Research under the 2010 Programme for Development of Higher Education, by the National Science Foundation under grant DMS 0604386 and by the Edgar Odell Lovett Fund at Rice University. B.M.G. is supported in part by the Russian Foundation for Basic Research grants 07-01-92215 CNRS(L) and 08-01-00105.

References

- [1] Abramov L.M., Rokhlin V.A. *The entropy of a skew product of measure-preserving transformations*, Vestnik Leningrad. Univ. 17 (1962), 5–13. Amer. Math. Soc. Transl. (Ser. 2), 48 (1965), 225–65.
- [2] Athreya J. *Quantitative recurrence and large deviations for Teichmüller flows*, Geom. Dedicata, 119 (2006), 121–140.
- [3] Avila A., Gouëzel S., Yoccoz J.-C. *Decay of correlations for Teichmüller flows*, Publ. Math. IHES, no. 104 (2006), 143–211.
- [4] Bufetov A.I. *Decay of Correlations for the Rauzy–Veech–Zorich Induction Map on the Space of Interval Exchange Transformations and the Central Limit Theorem for the Teichmüller Flow on the Moduli Space of Abelian Differentials*. J. Amer. Math. Soc., 19 (2006), no. 3, 579–623.
- [5] Bufetov A.I., Gurevich B.M. *Existence and Uniqueness of the Measure of Maximal Entropy for the Teichmüller Flow on the Moduli Space of Abelian Differentials* (Russian). Funk. Anal. Pril., 42 (2008), 275–277.
- [6] Buzzi J., Sarig O. *Uniqueness of equilibrium measures for countable Markov shifts and multi-dimensional piecewise expanding maps*, Ergodic Theory and Dynamical Systems 23 (2003), no. 5, 1383–1400.
- [7] Cornfeld I., Fomin S., Sinai Ya. *Ergodic theory.*, Springer Verlag, 1982.

- [8] Forni, G. Devitaion of ergodic averages for area-preserving flows on surfaces of higher genus. *Ann of Math.*(2), 155(2000), 1-103.
- [9] Gurevich B.M. *Shift entropy and Markov measures on the path space of a countable graph* (Russian). *Doklady Akad Nauk SSSR* 192 (1970), 963-965.
- [10] Gurevich B.M. *A variational characterization of one-dimensional countable state Gibbs random fields*. *Z. Wahr. verw Geb.*, 68 (1984), 205–242.
- [11] Gurevich B.M., Katok S. *Arithmetic codeding and entropy for the positive geodesic flows on the modular surface*. *Mosc. Math. J.*, 2001, v. 1, no. 4, p. 569–582.
- [12] Gurevich B.M., Savchenko S.V. *Thermodynamic formalism for countable symbolic Markov chains* (Russian). *Uspekhi Mat. Nauk*, 53, no.2 (1998), 3-106. Engl. Transl. in *Russian Math. Surv.*, 53, no.2 (1998), 245-344.
- [13] Kontsevich M., Zorich A. *Connected components of the moduli spaces of Abelian differentials with prescribed singularities*. *Invent. Math.* 153 (2003), no. 3, 631–678.
- [14] G.A. Margulis, *On some aspects of the theory of Anosov systems*. With a survey by Richard Sharp: *Periodic orbits of hyperbolic flows*. Springer Monographs in Mathematics. Springer-Verlag, Berlin, 2004.
- [15] Masur H. *Interval exchange transformations and measured foliations*. *Ann. of Math.* (2) 115 (1982), no. 1, 169–200.
- [16] Rauzy G. *Échanges d'intervalles et transformations induites*. *Acta Arith.* 34, (1979), no. 4, 315–328.
- [17] Rokhlin V.A. *Selected questions of metric theory of dynamical systems*, *Uspehi Matematicheskikh Nauk*, 4 (1949), no. 2(30), 57–128.
- [18] Rokhlin V.A. *Lectures on entropy theory of transformations with invariant measure*. *Uspekhi Matematicheskikh Nauk*, 22 (1967), no. 5, 3–56.
- [19] Sarig O. *Thermodynamic formalism for countable Markov shifts*. *Ergod. Theory Dynam. Systems* 19 (1999), no. 6, 1565–1593.

- [20] Savchenko S.V. *Special flows constructed by countable topological Markov chains*. Func. Anal. Appl., 32 (1998), 1, 40–53.
- [21] Veech W. *Gauss measures for transformations on the space of interval exchange maps*, Annals Math., 15 (1982), 201–242.
- [22] Veech W. *Interval exchange transformations*. J. Analyse Math., 33 (1978), 222–272.
- [23] Veech W. *The Teichmüller geodesic flow*. Annals of Mathematics (2) 124 (1986), no. 3, 441–530.
- [24] Viana M. *Stochastic dynamics of deterministic systems*, Brazilian Colloquium of Mathematics, IMPA.
- [25] Viana M. *Dynamics of interval exchange maps and Teichmüller flows*. Lecture notes of graduate courses taught at IMPA in 2005 and 2007.
- [26] Zorich A. *Finite Gauss measure on the space of interval exchange transformations. Lyapunov exponents*. Ann. Inst. Fourier (Grenoble), 46 (1996), no. 2, 325–370.
- [27] Zorich A. *Flat surfaces*. In: Frontiers in Number Theory, Physics, and Geometry, Vol. 1 (2006), 439–586.